

DESY 98-013

IFT 97/14

hep-ph/9806291

February 1998

Survey of recent data on photon structure functions and resolved photon processes

Maria Krawczyk^a and Andrzej Zembrzuski

Institute of Theoretical Physics, Warsaw University, Poland

^a and Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany

and

Magdalena Staszal

Institute of Experimental Physics, Warsaw University, Poland

Abstract

Present data on the partonic content of the photon are reviewed. The results on the unpolarized structure functions from DIS experiments and on large p_T jet production processes in $\gamma\gamma$ and γp collisions are discussed for both real and virtual photons. A few related topics like the QED structure functions of the photon and the structure function of the electron are also shortly discussed.

Contents

1	Introduction	1
2	Partonic content of the real photon	3
2.1	DIS _γ experiments	3
2.1.1	Theoretical description	4
2.1.2	Data on F_2^γ	6
2.2	Hadronic final states in the DIS _γ experiments	25
2.2.1	Modeling of the hadronic final state in $\gamma^*\gamma$ collision	25
2.2.2	Data on the hadronic final state	26
2.3	Large p_T processes in $\gamma\gamma$ and γp collision	38
2.3.1	General framework	39
2.3.2	Measurements of the resolved photon processes	40
2.3.3	Jet production in $\gamma\gamma$ collision	42
2.3.4	Jet production in the resolved γp scattering	51
3	Partonic content of the virtual photon	78
3.1	Theoretical framework	78
3.2	DIS _{γ*} for virtual photons	79
3.3	Jet production in resolved virtual photon(s) processes in $\gamma\gamma$ and γp collisions	80
4	Related topics	85
4.1	On leptonic structure functions of the photon	85
4.2	On the structure function of the electron	91
5	Summary and outlook	92
6	Appendix	94
6.1	Parton parametrizations for the real photon	94
6.2	Parton parametrizations for the virtual photon	95
7	References	96

1 Introduction

The concept of the hadronic (partonic) structure of photon is used in describing high energy photon - hadron interactions (for a general discussion see *e.g.* review articles [1]). There are two basic types of inclusive processes where the structure of photon is tested in existing experiments:

- the deep inelastic scattering (DIS_γ), $e\gamma \rightarrow e \text{ hadrons}$, where the structure functions of photon $F_{1,2,\dots}^\gamma$ are measured,
- the large p_T jet production in γp and $\gamma\gamma$ collisions (one of the so called resolved photon processes), where individual quark and gluon densities in the photon may be probed.

The early experiments of the first type were performed at PETRA and PEP e^+e^- colliders [2]¹. Final analyses of LEP 1 data on F_2^γ taken at CM energy $\sim M_Z$ appeared recently, new data are being collected and analyzed at e^+e^- LEP collider running at higher energies (CM energy: 130-136 and 161-172 GeV), and at KEK collider (energy ~ 60 GeV).² All together they cover a wide range of Q^2 from 0.24 to 390 GeV²; the range of the x_{Bj} variable extends from ~ 0.002 to 0.98.

The latter type of measurements, *i.e.* measurement of the jets in the resolved photon processes, started a decade later³. Data have been and are still being taken in photoproduction processes at the ep collider HERA (with the CM energy 300 GeV) and in $\gamma\gamma$ collisions at the above mentioned e^+e^- machines. They have just started to give first results on the light quark (the effective parton density) and gluonic content of the photon.

In this paper the status of recent measurements of the structure function of unpolarized photons for both real and virtual photons in “ DIS_γ ” experiments and in large p_T jet processes involving resolved photons is discussed⁴. In the presentation we focus on recent results (data collected in years 1990 to 1996)⁵, where qualitative change has appeared in both types of measurements. Final results based on few years’ runs at LEP 1 are being published and a summing - up can be done. On the other hand, impressive progress has been obtained in pinning down the individual parton contributions in resolved photon processes with large p_T jets at HERA. These two methods of studying the “structure” of photon are becoming recently even more closely related, as it turned out that in the reconstruction of F_2^γ by the unfolding procedure the detailed analysis of the final hadronic state, with the contribution due to resolved photon processes, has to be performed.

The discrepancies observed by many collaborations in the description of the hadronic final states in the DIS_γ experiments have enforced the advanced study of various aspects of hadron production in this experimental setup. We found it important to

¹The first measurement was done in 1981 by the PLUTO Collaboration [2].

²In principle the measurement could be performed also at the SLC collider (at energy ~ 91 GeV).

³First evidence for the resolved process was found in 1990 by the AMY collaboration at KEK [3].

⁴So far there is no data for the polarized photon; for recent discussion of the polarized structure functions of photons see *e.g.* [4, 5].

⁵The exception is made for the structure of the virtual photon and the electromagnetic structure function for the photon, where old experimental data are also reviewed.

collect these results together and in close relation to the large p_T jet production results, where problems with the proper description of the underlying events appear as well. We hope that this way of presentation may help to clarify the situation with the production of hadrons in photon induced processes.

A very comprehensive paper "A compilation of data on two-photon reactions leading to hadron final states" by D. Morgan, M. R. Pennington and M. R. Whalley [6] contains data published up to March '94. Beside the inclusive hadron production it contains also the data on the exclusive processes and on the total cross sections, which are beyond the scope of the present paper.

Our survey is based on the published data, with few very recent preliminary results presented at the 1997 conferences. The short descriptions of results obtained by the experimental group together with some representative figures as well as comments/conclusions quoted from original publications are given. We describe the data by the name of the collaboration in alphabetic order, the publication date and the reference listed at the end of the paper. Each of the different topics is introduced by a short review of the basic theoretical ideas and notation.

In Sec.2, data for the real photon are discussed. Sec.2.1 deals with DIS_γ results (structure function F_2^γ), Sec.2.2 with data on hadronic final states in DIS_γ experiments for the real photon and Sec.2.3 with large p_T jet production in $\gamma\gamma$ and γp collisions. In Sec.3 existing data for the virtual photon are presented. In Sec.4 related topics are shortly discussed, namely leptonic structure functions of the photon (Sec.4.1) and the structure function of the electron (Sec.4.2). Sec.5 contains the summary and outlook. In the Appendix, the existing parton parametrizations are listed.

2 Partonic content of the real photon

2.1 DIS $_{\gamma}$ experiments

In this section we consider the standard DIS $_{\gamma}$ measurements in unpolarized $e\gamma$ collisions based on the process (Fig. 1):

$$e\gamma \rightarrow e \text{ hadrons}, \quad (1)$$

at e^+e^- colliders (so called single - tagged events) [1]. Here the target is a real (in practice almost real) photon, coming from the initial electron (positron).




Figure 1: Deep inelastic scattering $e\gamma$ ($P^2 \approx 0$) $\rightarrow eX$. The photon probe has virtuality $q^2 = -Q^2$.

The flux of these real photons in present experimental setups can be approximated by the Weizsäcker - Williams formula (called also Equivalent Photon Approximation EPA) [7, 1].⁶ It corresponds to the typical soft bremsstrahlung spectrum.⁷ There is a possibility to introduce here the structure function of the electron, which would contain the convolution of the Weizsäcker - Williams flux for virtual photons and the structure function of γ^* (see discussion in Sec.3 and 4).

The cross section for process (1), where the real photon with four momentum p is probed by the virtual photon with four momentum $q = k - k'$ ($q^2 = -Q^2 < 0$), is given by:

$$\frac{d\sigma^{e\gamma \rightarrow eX}}{dx_{Bj}dy} = \frac{4\pi\alpha^2(2p \cdot k)}{Q^4} [(1-y)F_2^\gamma + x_{Bj}y^2F_1^\gamma] \quad (2)$$

$$= \frac{2\pi\alpha^2(2p \cdot k)}{Q^4} [(1 + (1-y)^2)F_2^\gamma - y^2F_L^\gamma], \quad (3)$$

with the relation among the transverse F_T^γ ($= F_1^\gamma$) and longitudinal F_L^γ structure functions: $F_2^\gamma = F_L^\gamma + 2x_{Bj}F_1^\gamma$. $k(k')$ are four momenta of initial (final) electron and

⁶Recent critical discussion of this approach for the $\gamma\gamma$ processes in e^+e^- collisions at LEP 2 can be found in [8].

⁷Note that at $e\gamma$ or $\gamma\gamma$ options planned at NL Colliders, where the real photons would be obtained in the Compton backscattering process, the corresponding flux is expected to be much harder [9].

the standard DIS variables are defined by

$$x_{Bj} = \frac{Q^2}{2p \cdot q}, \quad y = \frac{p \cdot q}{p \cdot k}. \quad (4)$$

Note that in the LAB frame y describes the scaled energy of the exchanged photon. In practice, the variable y is small and the cross section (2) is effectively saturated by F_2^γ . Note that in e^+e^- collisions there is always a small off-shellness of the photon-target, $p^2 = -P^2 \neq 0$, and there may appear in addition the third structure function, F_3^γ . It disappears however from the cross section in the single-tag measurements after integration over the azimuthal angle (since it enters the cross section as the term $F_3^\gamma \cos(2\phi)$). See Sec. 4.1 for the discussion on the additional structure functions.

2.1.1 Theoretical description

In the Quark Parton Model (QPM or PM) one assumes that the hadronic final state in Eq. (1) is due to the production of quark pairs: q_i and $\bar{q}_i$ ($i=1,2,\dots,N_f$ - number of flavours) with the fractional charge Q_i . The F_2^γ is obtained by the integration over the transverse momentum of the outgoing quark with respect to the target photon direction. The full (PM or the lowest order) expression for F_2^γ , keeping the terms with quark mass m_{q_i} , is given by the Bethe-Heitler formula [1]:

$$F_2^\gamma = \frac{\alpha}{\pi} N_c \sum_{i=1}^{N_f} Q_i^4 x_{Bj} [(-1 + 8x_{Bj}(1 - x_{Bj}) - x_{Bj}(1 - x_{Bj}) \frac{4m_{q_i}^2}{Q^2})\beta + [x_{Bj}^2 + (1 - x_{Bj})^2 + x_{Bj}(1 - 3x_{Bj}) \frac{4m_{q_i}^2}{Q^2} - x_{Bj}^2 \frac{8m_{q_i}^2}{Q^2}] \ln \frac{1 + \beta}{1 - \beta}], \quad (5)$$

where the quark velocity β is given by

$$\beta = \sqrt{1 - \frac{4m_{q_i}^2}{s}} = \sqrt{1 - \frac{4m_{q_i}^2 x_{Bj}}{Q^2(1 - x_{Bj})}}. \quad (6)$$

The energy of the $\gamma^*\gamma$ collision, s , is equal to the the square of the invariant mass of the hadronic system W^2 and is given by

$$s = W^2 = \frac{Q^2}{x_{Bj}}(1 - x_{Bj}). \quad (7)$$

In the limit of s well above threshold, *i.e.* for $\beta \approx 1$, which for fixed x_{Bj} (not too small and not too close to 1) corresponds to the Bjorken limit, one gets

$$\ln \frac{1 + \beta}{1 - \beta} = \ln \frac{(1 + \beta)^2}{1 - \beta^2} \approx \ln \frac{Q^2(1 - x_{Bj})}{m_{q_i}^2 x_{Bj}}. \quad (8)$$

The structure function F_2^γ can be approximated in such case by

$$F_2^\gamma = \frac{\alpha}{\pi} N_c \sum_{i=1}^{N_f} Q_i^4 x_{Bj} [[x_{Bj}^2 + (1 - x_{Bj})^2] \ln \frac{W^2}{m_{q_i}^2} + 8x_{Bj}(1 - x_{Bj}) - 1] \quad (9)$$

and it can be used to define the quark densities in the photon:

$$F_2^\gamma = x_{Bj} \sum_{i=1}^{2N_f} Q_i^2 q_i^\gamma(x_{Bj}, Q^2). \quad (10)$$

In the above formulae $N_c=3$ denotes the number of colors. In the last equation a (natural) assumption, that quark and antiquark distributions in photon are the same, has been introduced. Note that

- F_2^γ is calculable in the PM, in contrast to the structure function of hadrons, *e.g.* the nucleon structure function F_2^N .
- F_2^γ is proportional to α , the fine structure coupling constant ($\approx 1/137$). There is an overall logarithmic dependence on the energy scale squared W^2 (or Q^2 , see Eq. (7)). A large value of F_2^γ , or in other words a large quark density, is predicted at large x_{Bj} .
- F_2^γ is a sum of the quark densities with the factors Q_i^4 , where Q_i is fractional charge of quark, so the individual quark density is proportional to Q_i^2 (see Eqs. (9, 10)).

The longitudinal structure function F_L^γ is not zero in PM, in contrast to the corresponding function for hadrons, and it is scale invariant:

$$F_L^\gamma = \frac{\alpha}{\pi} N_c \sum_{i=1}^{N_f} Q_i^4 x_{Bj} [4x_{Bj}(1 - x_{Bj})]. \quad (11)$$

Note that also the third structure function is scale invariant:

$$F_3^\gamma = \frac{\alpha}{\pi} N_c \sum_{i=1}^{N_f} Q_i^4 x_{Bj} [-x_{Bj}^2]. \quad (12)$$

In the leading logarithmic approximation (LLA) the Callan-Gross relation $F_2^\gamma = 2x_{Bj}F_1^\gamma$ holds as in the case of hadrons.

In this approximation the PM formula for the quark density is given by

$$q_i^\gamma(x_{Bj}, Q^2)|_{PM}^{LL} = \frac{\alpha}{2\pi} N_c Q_i^2 [(x_{Bj}^2 + (1 - x_{Bj})^2) \ln \frac{Q^2}{\Lambda_{QCD}^2}]. \quad (13)$$

Since the LL contribution corresponds to the on-shell quarks one can treat x_{Bj} in the above formula as equal to the part of four momentum of the initial photon carried by the quark; this latter variable is usually denoted by x_γ (or simply x). Note also that in the above formula instead of a quark mass there appears the QCD scale Λ_{QCD} . Therefore this expression allows to describe all the light quark (light as compared to the scale W^2) contributions to F_2^γ in a universal way. Heavy quark contributions should be treated separately, according to the Eqs. (5, 6) [1].

An additional Q^2 dependence will appear in F_2^γ and in q_i^γ due to the QCD corrections, which can be described by the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi equations or by other techniques [1, 10, 11, 12, 13]. In this framework the gluonic content of photon, g^γ , appears as well.

The inhomogeneous DGLAP equations for the real photon can be represented in the following way (below q_i^γ is used for quarks and antiquarks):

$$\begin{aligned}\frac{\partial q_i^\gamma}{\partial \ln Q^2} &= \frac{\alpha}{2\pi} Q_i^2 P_{q\gamma} + \frac{\alpha_s}{2\pi} \int_x^1 \frac{dy}{y} [P_{qq}(\frac{x}{y}) q_i^\gamma(y) + P_{qg}(\frac{x}{y}) g^\gamma(y)] \\ \frac{\partial g^\gamma}{\partial \ln Q^2} &= 0 + \frac{\alpha_s}{2\pi} \int_x^1 \frac{dy}{y} [P_{gq}(\frac{x}{y}) \sum_{i=1}^{2N_f} q_i^\gamma(y) + P_{gg}(\frac{x}{y}) g^\gamma(y)]\end{aligned}\quad (14)$$

with the standard splitting functions P_{qq} , P_{qg} , P_{gq} , P_{gg} and in addition with the function

$$P_{q\gamma} = N_c [x^2 + (1-x)^2], \quad (15)$$

describing the splitting of the photon into quarks.

Note that it is possible to solve the above equations without the initial conditions [10], assuming the LL or NLL behaviour of the solution. Obtained in this way the, so called, asymptotic solutions, have a singular behaviour at small x [13]. Therefore in practice while solving the equations (14) the initial conditions have to be assumed from a model or taken from measurements at some (low) Q_0^2 scale [14]. On this basis the parton parametrizations for the photon can be constructed, see the Appendix for the details of existing parton parametrizations.

In pre-QCD times, the hadronic structure of the photon has been solely attributed to the vector meson component (ρ, ω, ϕ) in the (real) photon. Therefore for the matrix element of the photon between *e.g.* the nucleon states, the following representation by the corresponding matrix elements for the ρ current was assumed [15]:

$$\langle N | J_\mu | N \rangle = -\frac{m_\rho^2}{g_\rho} \frac{1}{q^2 - m_\rho^2} \langle N | \rho_\mu | N \rangle + \dots \quad (16)$$

That was the basic assumption of the Vector Dominance Model (VDM) or Generalized VDM (GVDM) if higher vector mesons states were included [15, 16]. This non-perturbative (or hadronic) contribution to the structure of the photon is present of course in the measured structure function F_2^γ , as well as in q^γ or g^γ extracted from the experimental data.

2.1.2 Data on F_2^γ

Here we discuss the latest results for F_2^γ for a real photon based on single-tagged events at e^+e^- colliders (*i.e.* DIS $_\gamma$ events, see Eq. (1) and Fig. 1). Note that in practice not x_{Bj} but the quantity x_{vis} is measured,

$$x_{vis} = \frac{Q^2}{Q^2 + W_{vis}^2}, \quad (17)$$

where W_{vis} is the invariant mass of the visible hadronic system. For the nonzero target-photon virtuality P^2 , the measured quantity is

$$x_{vis} = \frac{Q^2}{Q^2 + W_{vis}^2 + P^2}. \quad (18)$$

From a measured value for x_{vis} one can reconstruct x_{Bj} by an unfolding procedure.

Recently, in order to perform reliable unfolding, the events are grouped into classes of similar topology (see *e.g.* **DELPHI 96b**), which roughly coincide with the contributions due to QPM, VDM or RPC (RPC - Resolved Photon Contribution, where photons interact through their partonic contents).

In the early analyses the separation between these contributions (QPM with QCD corrections and VDM, basically) was obtained by introducing the parameter of the minimal transverse momentum of the produced hadrons, p_T^0 . The VDM (or hadronic) contribution should be negligible for the production of particles with p_T larger than p_T^0 , where, on the other hand, other contributions should be included. The FKP approach [17], where this kind of cutoff parameter is built in, was natural here. Note that in generation of two quark final state (in *e.g.* F2GEN event generators) two extreme types of angular dependence for final quarks are assumed: point-like (PL) with angular dependence as in the lepton pair production in real photon-photon collision and a “peripheral” one with the distribution corresponding to an exponential form of the p_T dependence, with a mean of 300 MeV (as if the photon interacted as a hadron)⁸. Nowadays the Monte Carlo generators HERWIG and PYTHIA, adapted for DIS_γ in 1995, can describe all these types of contributions with any of the existing parton parametrizations. It is no longer necessary to fit an empirical p_T^0 parameter to the data before unfolding.

It is worth noticing here, that at present energies (or Q^2 scales) light quark (u, d, s) and heavy quark distributions are treated differently. For c, b quarks the QPM formula (Eq. (5)) is applied (at least close to the threshold), while for the lighter quarks the QCD corrections are necessary. Note also that in the earlier measurements the c quark contribution was usually subtracted from the structure function F_2^γ .

Unfolding can be performed using the traditional method based on linear scale or using the new approach with the logarithmic scale, especially useful for extracting F_2^γ results in the small x_{Bj} region.

As we have already pointed out, the study of the hadronic final state became part of the measurements of the structure function for the photon at e^+e^- colliders. The details of the studies of the hadronic final state will be presented in the next section. Here we would only like to stress that some discrepancies have been found for certain distributions, like the pseudorapidity η of final hadrons distributions. This fact is included in the estimation of the uncertainty of the measured function F_2^γ .

The general features of F_2^γ , as far as x_{Bj} and $\log Q^2$ dependences are concerned, agree with the theoretical expectations, although the precision of the data does not allow in many cases to distinguish between existing parton parametrizations and/or clarify the small x_{Bj} behaviour of F_2^γ .

We start the presentation of the data from the LEP collider, then TRISTAN data are discussed. Collective figures of F_2^γ versus x_{Bj} and F_2^γ versus Q^2 are presented at the end of this section.

The sets of data are described by the name of the collaboration (in alphabetic order), the publication date, and reference listed at the end of the paper. In comments we quote statements from the original papers (for abbreviations used for the parton

⁸also a mixed ‘perimiss’ dependence was studied [24].

parametrizations see the Appendix).

DATA

•ALEPH 97a [18] (LEP 1)

Data on F_2^γ at LEP 1 energy for x_{Bj} from 0.002 to 0.9 and Q^2 between 6 and 44 GeV^2 were collected in the period 1991-94. An analysis of the hadronic final state was performed using the QPM+VMD model and the HERWIG simulations (with GRV, GS and LAC1 parton parametrizations) (see next section).




Figure 2: The unfolded F_2^γ/α for $\langle Q^2 \rangle = 8.9$ and 19.1 GeV^2 compared with predictions of GRV, LAC1, and SaS1D parametrizations.

The unfolded results for F_2^γ are presented in Fig. 2. F_2^γ averaged over x_{Bj} and Q^2 in two bins of Q^2 is given in the table:

$\langle Q^2 \rangle$ [GeV^2]	x_{Bj}	F_2^γ/α (stat. + syst.)
8.9	0.3 – 0.8	$0.36 \pm 0.16 \pm 0.06$
19.1	0.3 – 0.8	$0.44 \pm 0.08 \pm 0.02$

Comment: Problem with the proper description of hadronic energy flow as a function of pseudorapidity or azimuthal separation angle was found (see next section).

•ALEPH 97b [19] (LEP I)

Data taken from 1991 to 1995 at average Q^2 equal to 279 GeV^2 . The hadronic final state was also studied using four QPM+VDM models and HERWIG Monte Carlo program (with GS2, GRV LO, LAC1 and SaS1d parton parametrizations, see next section). The results for F_2^γ are presented in Figs. 3 and 4.

Comment: Problem with the proper description of the pseudorapidity distribution of final hadrons (see next section).




Figure 3: The structure function F_2^γ/α as a function of x_{Bj} (from [19]).




Figure 4: The Q^2 dependence of F_2^γ/α for the range $0.3 < x_{Bj} < 0.8$ (from [19]).

•DELPHI 96a [20] (LEP 1)

Data on F_2^γ were taken in the period 1991-93 for Q^2 between 4 and 30 GeV^2 and for x_{Bj} down to 0.003. The so called $F_2^{\gamma(QED)}$ was also measured and compared with QED prediction, also for nonzero averaged virtuality P^2 of the target photon (see Sec. 4.1 for details). Estimated target photon virtuality P^2 was used in the unfolding of F_2^γ ⁹. The TWOGAM event generator was used to simulate QPM events and another event generator was used to obtain the QCD correction (LL) to the point-like contribution for light quarks in the FKP approach. The GVDm and the point-like (FKP) contributions were studied, with $p_T^0 = 0.1$ and 0.5 GeV. Results for

⁹ It was found that although $\langle P^2 \rangle = 0.13 \text{ GeV}^2$, a fixed value of 0.04 GeV^2 fits the data better.

F_2^γ at $\langle Q^2 \rangle = 12 \text{ GeV}^2$ are presented in Fig. 5 and in the table below:

$\langle Q^2 \rangle$ [GeV ²]	x_{Bj}	F_2^γ/α (stat. + syst.)
12	0.003 – 0.080	$0.21 \pm 0.03 \pm 0.06$
	0.080 – 0.213	$0.41 \pm 0.04 \pm 0.05$
	0.213 – 0.428	$0.45 \pm 0.05 \pm 0.05$
	0.428 – 0.847	$0.45 \pm 0.11 \pm 0.10$




Figure 5: Unfolded F_2^γ/α for the light quarks (DELPHI and OPAL data). The curves show the sum of the GVMD model prediction multiplied by the threshold factor $1-x_{Bj}$ and the prediction of the FKP parametrization for the point-like part of F_2^γ , with different values of parameter p_T^0 : 0.1 GeV (upper line) and 0.5 GeV (lower line). The bottom curves show the GVDM contribution with different target masses (from [20]).

The averaged value of F_2^γ/α over the x_{Bj} range between 0.3 and 0.8 was extracted:

$\langle Q^2 \rangle$ [GeV ²]	$\langle F_2^\gamma/\alpha \rangle$
12	0.45 ± 0.08

For comparison with other measurements of the Q^2 dependence of the F_2^γ , see Fig.6.




Figure 6: F_2^γ/α averaged over x_{Bj} between 0.3 and 0.8. The curves show the FKP parametrization predictions for different values of the parameter p_T^0 (from [20]).

Study of the F_2^γ behaviour at $\langle Q^2 \rangle = 12 \text{ GeV}^2$ in the low x_{Bj} domain leads to following results (Fig. 7):

$\langle Q^2 \rangle$ [GeV ²]	x_{Bj}	F_2^γ/α (stat. + syst.)
12	0.003 – 0.046	$0.24 \pm 0.03 \pm 0.07$
	0.046 – 0.117	$0.41 \pm 0.05 \pm 0.08$
	0.117 – 0.350	$0.46 \pm 0.17 \pm 0.09$




Figure 7: Unfolded F_2^γ/α for $\langle Q^2 \rangle = 12 \text{ GeV}^2$ from the DELPHI experiment with the LO parametrizations LAC1, GS, DO, and GRV (from [20]).

Comment: No rise of F_2^γ at small x_{Bj} has been found. GRV and GS leading order parametrizations of the quark density in the photon are in agreement with data.

•DELPHI 96b [21] (LEP 1)

Measurement of the photon structure function F_2^γ (data collected in the years 1991-95) together with a study of the hadronic final state, in particular the resolved target photon contribution (see next section), is reported.

Two types of models for the extraction of the VDM contribution were used for the $< Q^2 > = 13 \text{ GeV}^2$ data (GVDM and "TPC/ 2γ "-type, for details see Ref.[21]). QPM, and RPC contribution with the GS2, SaS4 and GRV3 parton parametrizations were studied. The results for the $< Q^2 > = 106 \text{ GeV}^2$ were also obtained. The unfolding was done in a linear and logarithmic scale in x_{Bj} .

Fig. 8 shows the x_{Bj} dependence for both Q^2 samples.




Figure 8: Unfolded F_2^γ/α compared with QPM+GVDM+RPC (GS2) (solid line) and predictions of SaS4 (dashed line), GS2 (dashed-dotted) and GRV3 (dotted) parametrizations (from [21]).

The averaged value of F_2^γ/α for $0.3 < x_{Bj} < 0.8$ as a function of Q^2 was extracted:

$< Q^2 >$ [GeV ²]	$< F_2^\gamma/\alpha >$ (stat. + syst.)
13	$0.38 \pm 0.031 \pm 0.016$
106	$0.576 \pm 0.081 \pm 0.076$

Comment: *The importance of the final hadronic state topology was noticed and the study of a linear and logarithmic unfolding performed.*

•DELPHI 97 [22] (LEP 1, LEP 2)

This paper reports a recent study of F_2^γ in the Q^2 range between 3 and 150 GeV², based on data from the 1994-95 runs for energies around the Z^0 mass and from 1996 for energies between 161 and 172 GeV. An analysis of the hadronic final state is performed and compared with predictions of the TWOGAM Monte Carlo program, where QPM, VDM and RPC(GS2) parts are included (see next section).

Unfolded results for F_2^γ as a function of x_{Bj} and Q^2 are presented in Figs. 9a and 9b, respectively.

Comment: *The data for hadronic final state distributions, including energy flow versus pseudorapidity, agree with predictions of the TWOGAM simulations.*




Figure 9: a) F_2^γ/α versus x_{Bj} from the DELPHI experiment based on LEP 1 data for three values of $\langle Q^2 \rangle$: 6.5, 13, and 22 GeV^2 . The solid line corresponds to QPM+GVDM+RPC(GS2), dotted - QPM+GVDM and dashed - GVDM. b) F_2^γ/α averaged over x_{Bj} from 0.3 to 0.8. Results obtained from LEP1 and LEP2 data are shown together with results of a previous analysis of LEP1 data (from [22]).

•OPAL 94 [23] (LEP 1)

Measurement of F_2^γ at $\langle Q^2 \rangle = 5.9$ and 14.7 GeV^2 was performed using data from the period 1990-92. The VDM and pointlike contributions (in the FKP approach) are separated by the cutoff parameter p_T^0 found to be $0.27 \pm 0.10 \text{ GeV}$. Results compared with PLUTO and TPC/2 γ data are presented in Fig. 10.




Figure 10: Unfolded F_2^γ/α shown with previous measurements in other experiments at similar $\langle Q^2 \rangle$ values. The curves show contributions of VDM (dots), QCD-based model (dashes) and their sum (line) for a) $\langle Q^2 \rangle = 5.9 \text{ GeV}^2$ b) $\langle Q^2 \rangle = 14.7 \text{ GeV}^2$ (from [23]).

•OPAL 97a [24] (LEP 1)

The measurement of F_2^γ was done for $6 < Q^2 < 30 \text{ GeV}^2$ and $60 < Q^2 < 400 \text{ GeV}^2$, using the full sample of data at the Z peak (years 1990-95). The detailed analysis of the hadronic final states was performed and sizeable discrepancies with the expectations

were found especially at low x_{Bj} (see next section). The influence of the choice of different MC generators (HERWIG, PYTHIA) on F_2^γ was studied. The estimation of the P^2 values was done based on the SaS1 parametrization.

The results for the F_2^γ , presented in Fig. 11a,b,c, are as follows (the value of F_2^γ/α is given at the centre of the x_{Bj} bin):

$\langle Q^2 \rangle$ [GeV ²]	x_{Bj}	F_2^γ/α (stat. + syst.)
7.5	0.001 – 0.091	$0.28 \pm 0.02^{+0.03}_{-0.10}$
	0.091 – 0.283	$0.32 \pm 0.02^{+0.08}_{-0.13}$
	0.283 – 0.649	$0.38 \pm 0.04^{+0.06}_{-0.21}$
14.7	0.006 – 0.137	$0.38 \pm 0.01^{+0.06}_{-0.13}$
	0.137 – 0.324	$0.41 \pm 0.02^{+0.06}_{-0.03}$
	0.324 – 0.522	$0.41 \pm 0.03^{+0.08}_{-0.11}$
	0.522 – 0.836	$0.54 \pm 0.05^{+0.31}_{-0.13}$
135.0	0.100 – 0.300	$0.65 \pm 0.09^{+0.33}_{-0.06}$
	0.300 – 0.600	$0.73 \pm 0.08^{+0.04}_{-0.08}$
	0.600 – 0.800	$0.72 \pm 0.10^{+0.81}_{-0.07}$




Figure 11: F_2^γ/α data from the OPAL experiment for the number of flavours $N_f=4$. Curves in (a)-(d) show predictions of GRV and SaS 1D parametrizations (from [24]).

The measurement of F_2^γ/α as a function of Q^2 averaged over x_{Bj} range, $0.1 <$

$x_{Bj} < 0.6$, leads to following results (Fig. 11d):

$\langle Q^2 \rangle$ [GeV ²]	$\langle F_2^\gamma / \alpha \rangle$ (stat. + syst.)
7.5	$0.36 \pm 0.02^{+0.06}_{-0.12}$
14.7	$0.41 \pm 0.01^{+0.08}_{-0.04}$
135.0	$0.71 \pm 0.07^{+0.14}_{-0.05}$

and the slope $d(F_2^\gamma / \alpha) / d \ln Q^2$ is measured to be $0.13^{+0.06}_{-0.04}$.

Comment: *No correction for $P^2 \neq 0$ was made. Large discrepancies between the hadronic energy flow data and Monte Carlo simulations are observed at low x_{vis} , when the results are presented versus pseudorapidity or azimuthal angle (see next section).*

•OPAL 97b [25] (LEP 1)

Measurement of F_2^γ at LEP 1 was done for $\langle Q^2 \rangle = 1.86$ GeV² and $\langle Q^2 \rangle = 3.76$ GeV² as a function of x_{Bj} , reaching the lowest ever measured (center of bin) value: $x_{Bj} = 0.0025$. For a better sensitivity on the low x_{Bj} region the unfolding procedure was performed on a logarithmic scale. Final state topology was analysed as well (using the HERWIG, PYTHIA, F2GEN generators - both with the pointlike and the peripheral distributions). Discrepancy between the data (hadron energy flow) and results from the Monte Carlo generators, as well as between different models was found for low $x_{vis} < 0.05$ (see Sec.2.2).

Obtained values of F_2^γ / α are shown in Fig. 12 together with measurements from PLUTO and TPC/2 γ .

Comment: *“No correction for $P^2 \neq 0$ was made. GRV-HO is consistent with the low- x OPAL results in the lower Q^2 bin, but at higher Q^2 it underestimates the low x OPAL data. GRV-LO and SaS1D describe the unfolded results worse than GRV-HO. Shapes of measured F_2^γ are flat within the errors, but a small rise in the low x region is not excluded.”*




Figure 12: OPAL F_2^γ/α data (circles) as a function of x_{Bj} for $\langle Q^2 \rangle = 1.86 \text{ GeV}^2$ (a) and $\langle Q^2 \rangle = 3.76 \text{ GeV}^2$ (b). Also data from PLUTO (crosses) for $\langle Q^2 \rangle = 2.4, 4.3 \text{ GeV}^2$ and TPC/2 γ (squares) for $\langle Q^2 \rangle = 1.31, 2.83 \text{ GeV}^2$ are shown. The curves show predictions of GRV-HO (dots), GRV-LO (line), and SaS1D (dashed) parametrizations. The range of the x_{Bj} -bins of the OPAL results are marked at the tops of figures (from [25]).

•OPAL 97c [26] (LEP 2)

New data on F_2^γ from LEP 2 at the CM energies 161-172 GeV were collected in 1996. The $\langle Q^2 \rangle$ range lies between 9 and 59 GeV^2 . Also distribution of the final hadronic energy flow was studied (HERWIG, PYTHIA and F2GEN, see next section).

The unfolded F_2^γ/α as a function of x_{Bj} and Q^2 are presented in Figs 13, 14. A special study of the Q^2 dependence of F_2^γ is performed for different x_{Bj} ranges, see Fig. 15.

A fit to the new data at the energies 161-172 GeV and the previous OPAL set at 91 GeV for Q^2 from 7.5 to 135 GeV^2 , averaged over the x_{Bj} range of 0.1-0.6 (see Fig. 14), has the form:

$$F_2^\gamma(Q^2)/\alpha = (0.16 \pm 0.05_{-0.16}^{+0.17}) + (0.10 \pm 0.02_{-0.02}^{+0.05}) \ln(Q^2/\text{GeV}^2).$$

Comment: No correction for $P^2 \neq 0$ was made. Discrepancies are observed in the hadronic energy flow between the data and the HERWIG and PYTHIA simulations, especially at $x_{vis} < 0.1$. Accuracy of the data does not allow to see the expected different slope of F_2^γ versus $\log Q^2$ for different x_{Bj} ranges.




Figure 13: x_{Bj} -dependence of F_2^γ/α measured by OPAL for different value of $\langle Q^2 \rangle$: 9, 30, 14.5, 59, 11, and 41 GeV². For comparison the GRV LO (solid line) and SaS1D (dashed dotted line) parametrizations are shown; dotted line represents the hadronic component (VDM?) (HAD) and the dashed one the asymptotic solution (ASYM) (from [26]).




Figure 14: F_2^γ/α averaged over $0.1 < x_{Bj} < 0.6$, for the energy 91 GeV (open circles) and for the energies 161, 172 GeV (full circles). Predictions of the QCD calculation are shown by the lines: solid (GRV LO), dotted (SaS1D), and double-dotted (HO, based on GRV HO parametrization for light quarks); dashed line corresponds to the asymptotic solution (ASYM). The charm contribution is calculated separately (from [26]).




Figure 15: F_2^γ/α for the $0.1 < x_{Bj} < 0.6$ range from Fig. 14 has been subdivided into three smaller bins. The data are compared here only to HO predictions (from [26]).

•AMY 95 [27] (TRISTAN)

A high Q^2 measurement of the photon structure function F_2^γ was performed. Unfolded F_2^γ is presented in Fig. 16 (below are the corresponding numbers).

$\langle Q^2 \rangle$ [GeV ²]	$\langle x_{Bj} \rangle$	F_2^γ/α (stat. + syst.)
73	0.25	$0.65 \pm 0.08 \pm 0.06$
	0.50	$0.60 \pm 0.16 \pm 0.03$
	0.75	$0.65 \pm 0.11 \pm 0.08$
390	0.31	$0.94 \pm 0.23 \pm 0.10$
	0.69	$0.82 \pm 0.16 \pm 0.11$

The values of F_2^γ averaged over $0.3 < x_{Bj} < 0.8$ are as follows:

$\langle Q^2 \rangle$ [GeV ²]	$\langle F_2^\gamma/\alpha \rangle$
73	0.63 ± 0.07
135	0.85 ± 0.18

Comment: The observed x_{Bj} -behaviour of F_2^γ is consistent with the GRV parametrization and with the FKP one (fitted parameter p_T^0 is equal to 0.51 ± 0.39 GeV, if only AMY data are included, and 0.45 ± 0.07 GeV if all available, at that time, F_2^γ data are taken).




Figure 16: F_2^γ/α data from the AMY collaboration compared with a few parametrizations of the photon structure: FKP, QPM (quark parton model), VMD model (a,b), WHIT1, LAC1, DG (c,d), GRV, AFG (e,f). Upper (a,c,e) and lower (b,d,f) figures correspond to averaged $Q^2=73 \text{ GeV}^2$ and $Q^2=390 \text{ GeV}^2$, respectively (from [27]).

•AMY 97 [28] (TRISTAN)

The measurement of F_2^γ at $< Q^2 \geq 6.8 \text{ GeV}^2$ was performed; the results for F_2^γ versus x_{Bj} , together with earlier data, are presented in Fig. 17. The comparison with various parametrizations is given in Fig. 18.




Figure 17: The measured F_2^γ/α values compared with results of other experiments at $Q^2 \sim 4-9 \text{ GeV}^2$ (from [28]).




Figure 18: The F_2^γ/α values as a function of x_{Bj} . a) Comparison with the SaS1D, SaS2D and GRV predictions. Also shown are the heavy quark contributions from the SaS and GRV parametrizations. b) Comparison with WHIT1, WHIT4, WHIT6 and LAC1 predictions. The heavy quark contribution is shown for WHIT1 parametrization (from [28]).

Comment: “The x_{Bj} behaviour of the measured F_2^γ is consistent with the QCD-based predictions such as SaS, GRV-LO and WHIT parametrizations, but inconsistent with LAC1 predictions for x_{Bj} value around 0.07.”

•TOPAZ 94 [29] (TRISTAN)

The photon structure function F_2^γ has been measured for averaged Q^2 values from 5.1 to 338 GeV^2 (see table below). The one and two jet events in this sample are studied as well (see also Sec.2.2 and 2.3).

$< Q^2 >$ [GeV^2]	Q^2 [GeV^2]	$< x_{Bj} >$	x_{Bj}	F_2^γ/α (stat. + syst.)
5.1	3 – 10	0.043	$0.010 - 0.076$	$0.33 \pm 0.02 \pm 0.05$
		0.138	$0.076 - 0.20$	$0.29 \pm 0.03 \pm 0.03$
16	10 – 30	0.085	$0.02 - 0.15$	$0.60 \pm 0.08 \pm 0.06$
		0.24	$0.15 - 0.33$	$0.56 \pm 0.09 \pm 0.04$
		0.555	$0.33 - 0.78$	$0.46 \pm 0.15 \pm 0.06$
80	45 – 130	0.19	$0.06 - 0.32$	$0.68 \pm 0.26 \pm 0.05$
		0.455	$0.32 - 0.59$	$0.83 \pm 0.22 \pm 0.05$
		0.785	$0.59 - 0.98$	$0.53 \pm 0.21 \pm 0.05$




Figure 19: TOPAZ 94 data for F_2^γ/α versus x_{Bj} for $Q^2=5.1, 16, 80 \text{ GeV}^2$ and a comparison with parton parametrizations: a) FKP, QPM (parton model), VMD model and b) DG, LAC1, GRV (from [29]).

The results are presented in Fig. 19. To study the Q^2 dependence of F_2^γ , the averaged values of F_2^γ/α in the x_{Bj} -range from 0.3 to 0.8 were extracted at $\langle Q^2 \rangle = 16, 80$ and 338 GeV^2 (the numbers in parentheses are the results for light quarks alone):

$\langle Q^2 \rangle$ [GeV ²]	F_2^γ/α
16	0.47 ± 0.08 (0.38 $\pm$ 0.08)
80	0.70 ± 0.15 (0.49 $\pm$ 0.15)
338	1.07 ± 0.37 (0.72 $\pm$ 0.37)

Comment: The final hadronic state described by $QPM(c)+VDM+FKP(u,d,s)$ with $p_T^0=0.1, 0.5$ and 1 GeV , also by GRV, DG and LAC1 parton parametrizations, was studied.

For an overall comparison the collective figures of F_2^γ/α versus x_{Bj} (Fig. 20 [30]) and F_2^γ/α versus Q^2 (Figs. 21 [31], 22 [32]), containing also earlier data not discussed

here, are presented. In Fig. 20 the comparison with theoretical predictions for x_{Bj} dependence of F_2^γ based on SaS-1D (LO) and GRV NLO parton parametrizations are shown (note the logarithmic scale). For comparison with theoretical predictions of the Q^2 evolution for different x_{Bj} ranges see Fig. 22, and also Figs. 4, 6, 9b, 11d, 14, 15.

The effective parton density, as measured at HERA collider by H1 group (see **H1 97a** [33]) in the jet production from resolved photons, is compared with the F_2^γ data in Fig. 23 as well.




Figure 20: The photon structure function F_2^γ/α as a function of x_{Bj} in bins of Q^2 compared to the GRV NLO (solid line) and SaS-1D (LO) (dashed line) parametrizations of parton distributions in the photon (from [30]).




Figure 21: The photon structure function F_2^γ/α (data from various experiments) as a function of Q^2 (from [31]).




Figure 22: The Q^2 evolution of F_2^γ/α measured by OPAL ($0.1 < x_{Bj} < 0.6$), AMY ($0.3 < x_{Bj} < 0.8$), JADE ($0.1 < x_{Bj} < 1.$), DELPHI ($0.3 < x_{Bj} < 0.8$), TPC ($0.3 < x_{Bj} < 0.6$), TOPAZ ($0.3 < x_{Bj} < 0.8$). The lines are predictions of parametrizations averaged over different x_{Bj} ranges, HO calculation being a sum of GRV HO parametrization for light quarks and the charm contribution calculated independently from GRV; also the asymptotic solution is shown (ASYM, dashed line) (from [32]).




Figure 23: The scaling violation of the parton densities compared with the Q^2 dependence of the F_2^γ (averaged over the range $0.3 < x_{Bj} < 0.8$) (from [33]).

2.2 Hadronic final states in the DIS_γ experiments

Although a detailed analysis of hadronic final states, including those involving large p_T particles/jets, has not been the main aim of DIS_γ experiments, nowadays it has proved crucial in extracting the unfolded F_2^γ . The obtained experimental results, and problems that appear while describing hadronic energy flow, transverse energy, pseudorapidity and other distributions within QCD, deserve close attention and a separate treatment - this section is devoted to this subject.

It is worth noticing that as F_2^γ corresponds to the total cross section for hadron production in $\gamma^*\gamma$ collisions, all the regions: $Q^2 \ll p_T^2$, $Q^2 \sim p_T^2$ and $Q^2 \gg p_T^2$ contribute, with the bulk of the cross section being due to the soft production, *i.e.* with a relatively small p_T . In the next section dedicated analyses of the large p_T jet production in real $\gamma\gamma$ and γp processes will be presented. Some of the problems that will appear there are common with these presented here.

2.2.1 Modeling of the hadronic final state in $\gamma^*\gamma$ collision

In this section the $\gamma^*\gamma$ collisions resulting in the hadronic state will be discussed. As we have already mentioned in Sec.2.1.2, the QPM, VDM and RPC contributions are introduced to model the hadronic final state. In recent analyses the MC generators HERWIG and PYTHIA are being used with chosen parton parametrizations. In the OPAL analysis, the generator F2GEN (based on TWOGEN used in older analyses), with assumed two quarks in the final state is used in addition, for the comparison and systematic checks. In this generator one can easily put a specific assumption on the final states. For example different angular distributions may be assumed in the $\gamma^*\gamma$ CM system: pointlike (PL), *i.e.* as for lepton pair from two real photons, peripheral with an exponential distribution of the transverse momentum, or the “perimiss” combination [24]. In a recent DELPHI analysis, the TWOGEN generator (“fixed” in 1993) was successfully used, with QPM, VDM and RPC contributions included [21, 22].

As far as data on global hadronic variables like W_{vis} , Q^2 , M_{ij} (invariant mass of two jets) and other distributions are concerned, there is a fairly good description of the data by existing MC generators. The problems arise for the transverse energy out of the tag plane (the plane defined by the initial and tagged electrons) or for the energy flow per event as a function of pseudorapidity, $\eta = -\ln(\tan(\theta/2))$, where the polar angle is measured from the direction of the target photon. The discrepancies are very pronounced especially for small x_{Bj} .

The first observation of the disagreement between the data and MC models was made by the OPAL collaboration [24]. “The serious discrepancies between the data and any of the available Monte Carlo models are seen both within the central region of the detector ($|\eta| < 2.3$), where the energy flow is well measured, and in the forward region, where the energy can only be sampled.” It is clear that the unfolding of F_2^γ “will have large errors as long as the energy flow from different models remains in clear disagreement with the energy flow in the data, in particular in the region of $x_{vis} < 0.1$ and $Q^2 < 30 \text{ GeV}^2$ ” (from [24]). A similar effect has been seen by now by other groups.

The method used recently to improve the agreement with the data is the so called “HERWIG + power law p_t ” generator, where the additional power law spectrum of

the form $dk_T^2/(k_T^2 + k_0^2)$ was introduced (it seems to be needed at HERA (ZEUS) as well, see next section).

One should be aware that in hadronic final state in DIS_γ experiments two types of large scale may appear: Q^2 and p_T^2 , describing the transverse momentum of the final hadrons/jets. The bulk of the data corresponds to events with not very large p_T ; if for these events the relation $Q^2 \gg p_T^2 \gg P^2$ holds, the interpretation in terms of the photon interaction between one direct (γ^*) and one resolved real (γ) photon may be introduced. Then it is not clear what scale should be used in the parton density for a real photon $f^\gamma(x, \tilde{Q}^2)$: p_T or Q^2 . Moreover the processes corresponding to $Q^2 \ll p_T^2$ should be treated as being resolved from the point of view of both: a real and a virtual photon.

Note, that in recent analyses the dependence on the number of jets in the final state is studied.

2.2.2 Data on the hadronic final state

The analyses of the hadronic final state accompanying the measurements of F_2^γ discussed in Sec.2.1.2 as well as the results from independent analyses of large p_T hadron production in single tagged events are presented below.

For the two photons involved, we introduce the following notation for their squared (positive) virtualities: $P_1^2 = Q^2$ and $P_2^2 = P^2$ (with $P_1^2 \geq P_2^2$), where the corresponding variables used in the discussed DIS scattering are also given.

DATA

•ALEPH 97a [18] (LEP 1)

Hadronic energy flow in the azimuthal angle separation ϕ_{sep} and pseudorapidity η was measured for Q^2 between 6 and 44 GeV^2 . The results are presented in Fig. 24.




Figure 24: Energy flow as a function of ϕ_{sep} (a) and η (b). Predictions: QPM+VDM (solid line) and HERWIG generator with GRV (dashed line) (from [18]).

Comment: "Discrepancy with MC models (QPM + VDM and HERWIG (with parton parametrizations GS, GRV, LAC1)) when energy flow is plotted as a function of rapidity and azimuthal separation."

•ALEPH 97b [19] (LEP 1)

Data taken from 1991 to 1995 for the average $Q^2=279 \text{ GeV}^2$. Four different QPM+VDM

models were used and in addition the HERWIG 5.9 generator with the GS, GRV, LAC1 and SaS1d parton parametrizations.

The pseudorapidity distribution compared with the model predictions is presented in Fig. 25. A disagreement is found, similar to the one observed previously by ALEPH at lower Q^2 (**ALEPH 97a**), but now at the negative pseudorapidity.




Figure 25: Pseudorapidity of all charged tracks and neutral calorimeter objects. Histograms are predictions of three of the models (dotted line from HERWIG, others - QPM+VDM type) (from [19]).

Comment: “The pseudorapidity distribution can not be described by the models as observed previously at lower Q^2 , note however that the excess is now in the opposite hemisphere than at lower Q^2 .”

•ALEPH 97c [34] (LEP 1)

Single tagged events collected in the years 1992-1994 at the average $Q^2=14.2 \text{ GeV}^2$. A dedicated study of the final hadronic state assuming the QPM+VDM for u, d and c quarks and using HERWIG 5.9 generator (with GRV parametrizations). Also the modified HERWIG generator was used which corresponds to the “HERWIG + power law p_t ”, with $k_{t0} = 0.66 \text{ GeV}$ (see also **OPAL 97e** where this model was introduced). Results are presented in Figs. 26, 27, 28. The approach “HERWIG + power law p_t ” leads everywhere to better description of the data.




Figure 26: The energy (out of plane of the tag and the beam) distribution (from [34]).

In order to pin down the source of observed discrepancy the analysis of the number of cone jets was performed in the final hadronic state (see **OPAL 97d** for the first

analysis of this type). HERWIG model where resolved photon processes are included should in principle give larger number of two jet events than QPM+VDM model, having additional production mechanisms. On the contrary, it gives less (see Fig. 28).

Also the energy not assigned to jets in two jet events, $E_2^{non-jet}$, was studied. To check the presence of the single and double resolved photon events, x_γ and x_{tag} distributions were measured. The “HERWIG + power law p_t ” approach was used successfully to describe these data as well.




Figure 27: The energy flow versus pseudorapidity of the final hadrons (from [34]).




Figure 28: (from [34]).

Comment: ”“HERWIG + power law p_t ” is better in modeling the region of large $E_{t,out}$ and the peak at positive rapidity”.

•**DELPHI 95 [35] (LEP 1)**

First evidence of hard scattering process in the single-tagged $e\gamma$ collision in the data from the 1991-1992 run is reported. The values of E_T of observed jets were larger than 1.5 GeV, while the magnitude of the mass of virtual photon was equal to $\langle P_1^2 \rangle \approx 0.06 \text{ GeV}^2$. This corresponds to the standard resolved (almost real) photon process, from the point of view of both photons.

Comment: This is not typical DIS experiment since the photon probe is almost real. The analysis is not typical, either, for the standard large p_T jet study. "The data are consistent with the predictions for quark and gluon density functions in the GS parametrization. The sum of the contributions: VDM + QPM (Quark Parton Model) + [QCD - RPC(Resolved Photon Contribution)] is needed in order to describe the data; the DO and LAC3 parametrizations do not adequately describe the data."

•**DELPHI 96b [21] (LEP 1)**

The results for the averaged $Q^2=13$ and 106 GeV^2 were considered and compared with TWOAM generator. The QPM, GVDM contributions and, for the first time at DELPHI, RPC with GS2 parton parametrization were included.

Comment: The resolved photon contributions are needed to obtain the description of the data. "...the correct unfolding procedure leads to the pointlike plus hadronic part of F_2^γ in a shape that is inconsistent with QPM (FKP) + GVDM in the region of low- x_{Bj} ."

•**DELPHI 97 [22] (LEP1, LEP2)**

Analysis of hadronic final state in the F_2^γ measurement (see Sec.2.1.2) was performed for energies around the Z^0 mass and for 161-172 GeV (1996 run). The Q^2 ranges: 6, 13, 22 and 17, 34, 63 GeV^2 were studied, respectively. Hadronic final state topology with events containing jets was studied using TWOAM generator. For results see Fig. 29 for LEP2 data and Fig. 30 for event energy flow at LEP1 and LEP2.

Comment: "All variables are in good agreement with TWOAM predictions". Note that the p_T range of jets may be of order of $\langle Q^2 \rangle$ for small Q^2 samples.




Figure 29: The comparison of data and the MC prediction for $\langle Q^2 \rangle = 34 \text{ GeV}^2$ (LEP2). a) Energy of tagged particle; b) The invariant mass; c) The number of jets; d) The jet angle with respect to the tagged particle; e) p_T^{jet} for jets in the same hemisphere as the tagged particle; f) p_T^{jet} for jets in the opposite hemisphere. Curves show the MC predictions: GVDM+QPM+RPC (GS2) (solid line), GVDM+QPM (dots), GVDM (dashes) (from [22]).




Figure 30: The event energy flow as a function of the pseudorapidity. The notations as in Fig. 29 (from [22]).

•OPAL 94 [23] (LEP 1)

Data were collected in 1990-92 for averaged $Q^2 = 5.9$ and 14.7 GeV^2 . Early analysis

of the final hadronic state (the TWOGEN generator with the contribution based on QPM, VDM and on the FKP approach to the QCD contribution) was performed. For the estimation of the p_T^0 see Sec.2.1.2.

•OPAL 97a [24] (LEP 1)

The hadronic final state was analysed in the measurement of the F_2^γ with one photon highly virtual (Q^2 between 6-30 GeV² - *low Q^2 sample* and between 60-400 GeV² - *high Q^2 sample*), the other being almost real.

For generating the hadronic final state MC programs HERWIG, PYTHIA, and for comparison F2GEN were used with GRV and SaS1D parton parametrizations.

The energy ($E_{t,out}$) transverse to the plane defined by the beam axis and the tag direction, and other quantities for the *low Q^2 sample* are presented in Fig. 31. The discrepancy found for the E_{vis} and $E_{t,out}$ distributions in this sample (Figs. 31 b and d, respectively) is absent in the *high Q^2 events* (not presented).




Figure 31: Comparison of data event quantities in the low- Q^2 sample with HERWIG, PYTHIA and F2GEN (PL) Monte Carlo simulations. a) the distribution of the visible invariant mass; b) the total visible energy of the event; c) the transverse momentum of the event in the tag plane; d) the energy out of the tag plane (from [24]).

The hadronic energy $E_{t,out}$ distributions in the three x_{Bj} bins are shown in Fig. 32. The failure of the models in the low Q^2 region is most visible at low x_{Bj} .

To establish the source of the discrepancy the energy flow per event in the *low Q^2 sample* was also studied as a function of the pseudorapidity (see Fig. 33). The distribution of pseudorapidity η for the *low Q^2 sample*, corrected for the experimental effects, is shown in Fig. 34.




Figure 32: The energy transverse to the tag plane for three x_{vis} bins for the low Q^2 sample. MC generators as in Fig. 31 (from [24]).




Figure 33: The hadronic energy flow per event as a function of the pseudorapidity η for the data and various MC simulations, in various ranges of x_{vis} and θ_{tag} for the low Q^2 sample (from [24]).




Figure 34: The measured energy flow in the low- Q^2 sample corrected for the detector inefficiencies, as a function of the pseudorapidity η , compared to the values generated by various MC models (from [24]).

Comment: None of the generators represents the final state accurately; $E_{vis}, E_{t,out}$ - distributions as well as the hadronic flow per event show a clear discrepancy. The failure of the models in the low Q^2 region is most marked at low x_{Bj} . The differences between MC models and data in the low Q^2 region are apparent, when the energy flow per event is plotted as a function of the pseudorapidity and the azimuthal angle (not shown). “Particular attention will need to be given to the angular distribution of partons in $\gamma^*\gamma$ system”.

The relation between p_T^2 and Q^2 in the low Q^2 sample may indicate a need to take into account the structure of the virtual photon.

•OPAL 97b [25] (LEP 1)

The measurement of F_2^γ and the modeling of the $\gamma^*\gamma$ fragmentation into hadrons at low Q^2 region (1.1 to 6.6 GeV^2) and very small x_{Bj} - bins from 0.0025 to 0.2 is reported. The hadronic energy flow as a function of η for three x_{Bj} regions is plotted in Fig. 35.

Comment: Differences in the energy flow distributions versus the pseudorapidity (Fig. 35) and in the summed energy transverse to the tag plane are found for $x_{vis} < 0.05$ between the data and MC models (HERWIG, PYTHIA, F2GEN).

•OPAL 97d [36] (LEP 1)

An analysis of the hadronic final state was done, in which the discrepancies between the data and predictions, reported in **OPAL 97a** [24], were examined from the point

of view of the number of produced jets. The data for $Q^2 \approx 6 - 30 \text{ GeV}^2$ taken in the years 1994-95 were compared to the results of the HERWIG, PYTHIA and F2GEN generators.




Figure 35: The measured hadronic energy flow as a function of the pseudorapidity: a comparison of data with various MC predictions for three bins in x_{vis} and two bins in θ_{tag} (from [25]).

The numbers of events for different number of jets, divided by the sum of all events, are presented in Fig. 36.




Figure 36: The fraction of events with 0-3 jets. The points show the data with the background subtracted, with statistical errors; histograms obtained with HERWIG, PYTHIA and F2GEN generators (from [36]).

The results of further studies of the energy flow versus the pseudorapidity η for events with different number of jets are shown in Fig. 37.




Figure 37: The hadronic energy flow per event in bins of the pseudorapidity η . Plots a), b), c) and d) show the average energy flow per event for summed events, and for events with 0, 1 and 2 jets, respectively. The samples are represented as in Fig. 36 (from [36]).

The number of events versus the energy transverse to the tag plane, obtained by the different generators and observed in the experiment, are compared in the Fig. 38.




Figure 38: The number of a) 1-jet and b) 2-jet events as a function of the jet transverse energy, E_T^{jet} . The HERWIG, PYTHIA and F2GEN events are normalized to the a) 1-jet and b) 2-jet data events (from [36]).

Comment: “There is a marked difference between the data and the Monte Carlo samples in the number of events with 2 jets”. “While the F2GEN sample, generated using the pointlike approximation, overestimates the 2 jet rate by a factor 2.4, HERWIG and PYTHIA are too low by factors of 2.4 and 3.6 respectively.”

Disagreement between the data and each of generators for the hadronic energy flow versus the rapidity is seen for all types of events: 0, 1 and 2-jet.

“All of the Monte Carlo samples model well the $E_{t,out}$ distributions for events with 1 jet, but PYTHIA underestimates the $E_{t,out}$ of the events with 2 jets.”

•OPAL 97e [37] (LEP 1)

An analysis of OPAL data taken in 1993-95 was performed at $\langle Q^2 \rangle = 7.5$ and 14.5 GeV^2 . The possible improvement of the description of the transverse energy flow in DIS_γ experiment by introduction of the additional intrinsic k_t smearing, influencing the angular distributions of the hadronic final state in the HERWIG, PYTHIA and ARIADNE generators was studied for the first time.

In Fig. 39 the transverse energy out of the tag plane is presented together with the results of the simulation by PYTHIA and HERWIG generators. The energy flow as a function of η was also studied for different x bins and Q^2 values (not shown).




Figure 39: The transverse energy out of the tag plane compared to PYTHIA (left) and HERWIG (right) simulations (from [37]).

Comment: “Both the $E_{t,out}$ and the energy flow per event are greatly improved with the inclusion of the power-like distributions of the intrinsic k_t , with the exception of the peak in the energy flow at low x_{vis} and $Q^2 = 14.7 \text{ GeV}^2$, which still falls short of the data.”

•TOPAZ 94 [29] (TRISTAN)

The jet production (one and two jets) has been studied in the deep inelastic $e\gamma$ scattering ($3.0 \text{ GeV}^2 < Q^2 < 30 \text{ GeV}^2$), see also Sec.2.1.2. Events with the transverse momentum of jets between 2 and 8 GeV and for the $|\eta| < 0.7$ were studied using the jet cone algorithm with $R=1$ (see next section for the definition of the jets).

The point-like and hadron-like configurations resulting in the different final-state topologies were studied.

Comment: *The two high p_T jet events are consistent with the point-like perturbative part. "In the one - jet sample an excess over the point like component is observed, which is direct experimental evidence for the existence of the hadron-like component in DIS $e\gamma$ ".*

2.3 Large p_T processes in $\gamma\gamma$ and γp collision

The structure function F_2^γ considered in Sec.2.1 is sensitive mainly to the combination of quark densities, and moreover, due to strong dependence on the charge, to the up -type quark distributions. Therefore it is of great importance to have an additional and very different from DIS_γ -experiments source of information. Hard processes involving resolved real photons provide such a source, with the cross section being a combination of the different parton densities convoluted with the cross sections for basic partonic subprocesses (see below). The role of these processes in determining *e.g.* a gluon density in the photon is unique indeed.

As we have already mentioned, the jet production in $\gamma\gamma$ collision leads to the complementary information on the photon structure to that coming from the F_2^γ ; moreover, the study of the hadronic final state has become recently a standard part of the structure function analyses performed at LEP (see discussion in Secs. 2.1 and 2.2).

In this section we focus on the large p_T jet production in the resolved $\gamma\gamma$ and γp collisions, where photon may interact through its partons. The main goal here is to extract the individual parton density in the real photon ¹⁰. Beside the jet production also the inclusive one-particle production is sensitive to the partonic content of photon, but it depends on the additional fragmentation functions, and will not be discussed here ¹¹. Also, other hard processes with resolved photon(s), like the Drell-Yan pair production may be used in pinning down the parton distributions in the photon. The lack of data reflects much lower rates for these processes.

The production cross section of jets, which are the hadronic representations of the hard partons produced in the basic subprocess, does depend however on the applied jet definition. Although this introduces additional uncertainty in the description of data, relatively high rate for these events makes large p_T jet production, apart from the F_2^γ , the basic source of information on the partonic structure of γ .

In the previous section we have discussed single tagged events with (positive) mass of the photons radiated by the tagged electron larger or much larger than 1 GeV (with exception of one experiment **DELPHI 95**). In this section the initial photons may be considered real, as the transverse momenta of the observed jets are chosen to be much larger than the median mass of the photon(s).

At present, the $\gamma\gamma$ collisions arising in e^+e^- colliders in no-tag ¹² conditions correspond to the $\gamma\gamma$ events where (both) real photons can be described by the Weizsäcker - Williams energy spectra. In the OPAL experiment, the squared target photon masses were estimated to be $\langle P_{1,2}^2 \rangle = 0.06 \text{ GeV}^2$, in the ALEPH experiment around 0.23 GeV^2 . In the γp scattering at ep collider HERA similar Weizsäcker - Williams spectrum describes the flux of photons coming from the electron (at HERA at present, *i.e.* in 1996/7, the positron) - here $P^2 < 4 \text{ GeV}^2$ (but with the median 0.001 GeV^2) or

¹⁰Presently most of the data deal with the almost real photon; for the jet production in processes with virtual photon, see next section.

¹¹with the exception of the newest charged particle production measurements by H1 group, where the gluon content has been derived.

¹²The name "untagged" is also used. In some cases antitagged or single - tagged events in the large p_T jet production are discussed in $\gamma\gamma$ and γp collisions as well (see below).

below 0.01-0.02 GeV², if the dedicated detector is used.

2.3.1 General framework

Resolved photon processes with large p_T jets can be characterized by the scale of hardness $\tilde{Q}^2$ which is usually provided by the p_T of the final jets, *i.e.* $\tilde{Q}^2 \sim p_T^2$. (We introduce here the notation $\tilde{Q}^2$ in order to distinguish it from the DIS scale Q^2 , which is equal to the virtuality of the photon probe.) In order to resolve both photons the scale $\tilde{Q}^2$ should be much larger than P^2 – the virtuality of the most virtual photon in the process, and moreover much larger than Λ_{QCD}^2 in order to apply the perturbative QCD. (Events where the masses fulfill the relation $\Lambda_{QCD}^2 \ll P^2 \ll \tilde{Q}^2$ are discussed in the next section, where the concept and data on the structure of virtual photon will be introduced.)

In this section initial photon(s) are considered to be real. Assuming the factorization between the hard subprocess and parton densities, the generic LO cross section for the two jet production in γp collision (Fig. 40a) is given by

$$d\sigma^{\gamma p \rightarrow jet_1 jet_2 X} = \sum_{i,j} \int \int dx_\gamma dx_p f_{i/\gamma}(x_\gamma, \tilde{Q}^2) f_{j/p}(x_p, \tilde{Q}^2) \hat{\sigma}^{ij \rightarrow jet_1 jet_2}, \quad (19)$$

where $f_{i/\gamma}(x_\gamma, \tilde{Q}^2)$ describes the probability in the LL approximation of finding a parton of a type "i" in the photon. The $x_\gamma(x_p)$ variable is by definition the part of the four momentum of the photon (proton) carried by its parton. Scale $\tilde{Q}^2$ is taken usually to be equal to the so called factorization scale. Note that the precise definition of the factorization scale can be given in the QCD calculation going beyond the LO approach.

For the two-jet production in $\gamma\gamma$ collision, $\gamma\gamma \rightarrow jet_1 jet_2 X$ (Fig. 40b) a similar formula holds:

$$d\sigma^{\gamma\gamma \rightarrow jet_1 jet_2 X} = \sum_{i,j} \int \int dx_\gamma^1 dx_\gamma^2 f_{i/\gamma}(x_\gamma^1, \tilde{Q}^2) f_{j/\gamma}(x_\gamma^2, \tilde{Q}^2) \hat{\sigma}^{ij \rightarrow jet_1 jet_2}. \quad (20)$$




Figure 40: Inclusive large p_T jet production in the resolved γp collision (a) and in the (double) resolved $\gamma\gamma$ collision (b).

Note that, contrary to the DIS _{γ} experiments, in the resolved photon processes there is no one basic observable, analogous to the F_2^γ . Usually the p_T or E_T and pseudorapidity η distributions are compared with the QCD prediction using a specific

parton parametrization ¹³ and the agreement justifies *a posteriori* the correctness of the applied parton parametrization and the QCD calculation. Only in few cases the parton distribution is extracted from data (see below).

It is worth noticing the difference between the variables describing the produced parton and the corresponding quantities for the jet, representing the considered parton on the hadronic level. The unfolding procedure is needed in order to obtain the "true" partonic variables, *e.g.* transverse momentum of the parton (p_t) from transverse momentum or energy of the jet (p_T or E_T).

2.3.2 Measurements of the resolved photon processes

For the jet production processes where at the Born level (*i.e.* in the LO) two jets appear, the measurements of basic distributions in E_T or the pseudorapidity η are performed for single or double jet (dijets) events as a rule.

For comparison with QCD not only the above distributions are important but also the study of the structure of the jet (jet profile or transverse energy flow around the jet axis) and the energy of the underlying event, or the hadronic activity outside the jets where effects due to accompanying remnant jets may be seen.

For more detailed study of properties of the resolved photon processes the separation of the direct photon events, where the photon participates directly in the hard subprocess, $x_\gamma \sim 1$, and resolved photon events is needed. Strictly speaking this separation holds only in the LO approach. In the two jet events, x_p and x_γ (or $x_\gamma^{1,2}$) distributions, sensitive to parton densities, can be reconstructed, see *e.g.* **ZEUS 95b, 97b**. The angular distribution (*e.g.* in $\gamma\gamma$ CM system) $d\sigma/d\cos\theta^*$, on the other hand, is not sensitive to these ingredients. So this measurement may help to verify the expectation of different angular distributions which correspond to direct and resolved photon contributions (*i.e.* due to different subprocesses), see **ZEUS 96a** for the first results.

The theoretical predictions based on the LO [94] - [96] or NLO(HO) QCD calculations are available for the inclusive one and two jet production for e^+e^- and ep experiments [97] - [110]. The main MC generators used by the experimental groups for these kinds of processes are PYTHIA, HERWIG and PHOJET (with or without the multiple parton interaction). They allow to study different types of distributions of the final state hadrons.

The following two jet definitions are used in the analyses reported below. The jet cone algorithm with the fixed value of the cone variable R , defined as $R = \sqrt{(\delta\phi)^2 + (\delta\eta)^2}$, with $\delta\phi$ ($\delta\eta$) describing the differences between the cone (jet) axis and the particle direction in the pseudorapidity and azimuthal angle, is used in PUCCELL and EUCELL generators. The second one corresponds to the k_T -cluster algorithm on which the KTCLUS generator is built up. Beside R also R_{sep} is included in some analyses, which corresponds to the additional separation between partons - note the $R_{sep} = 2 R$ means no restriction. (The discussion of jet definitions can be found in [110]).

¹³See the collection presented in Appendix.

The basic distributions, *e.g.* the jet E_T distribution, are in general in agreement with the expectation for both single and double jet production. In the pseudorapidity dependence the discrepancy is observed both for the jet rates and for the transverse energy flow around the jet axis, especially for small E_T and a small x_γ . This is taken as a hint that the multiple scattering may be important in the photon induced processes. Since the direct events should be free from such multiple interactions, the $\gamma\gamma$ and γp collisions may offer a unique laboratory to study this problem. The strong dependence on the proper choice of the jet definition and the parameters like the cone size R , is observed recently in the jet production at HERA. There is a possible relation of this effect to the problem of describing the underlying events observed both in e^+e^- and in ep collisions.

2.3.3 Jet production in $\gamma\gamma$ collision

Here the results for the jet production in $\gamma\gamma$ collision, where *one* or *two* resolved photons may interact, will be presented.

As in previous section we will denote the squared (positive) virtualities of two involved photons by P_1^2 and P_2^2 . In e^+e^- machine parton momentum fractions - variables $x_\gamma^{1,2} = x_\gamma^\pm$ - are determined from the two final jets with the highest E_T according to the formulae:

$$x_\gamma^\pm = \frac{\Sigma_{jets}(E \pm p_z)}{\Sigma_{hadrons}(E \pm p_z)}. \quad (21)$$

DATA

•DELPHI 94 [38] (LEP 1)

Data were collected in the period 1990-92 in antitagging conditions ($P_{max}^2 \sim 0.12 \text{ GeV}^2$). Jets defined according to the Lund cluster algorithm with p_T greater than 1.75 GeV were observed.

Results for the p_T of the jet distributions are shown in Fig. 41 (p_T^0 has been introduced to separate the resolved photon contribution calculated using the FPK approach with DG and DO parametrizations).




Figure 41: Distribution of jet transverse momentum compared with MC predictions (TWOAM generator). Dark area - QPM only; dots - QPM + VDM; full line - QPM + VDM + DG ($p_T^0 = 1.45 \text{ GeV/c}$); dashed line - QPM + VDM + DO ($p_T^0 = 1.22 \text{ GeV/c}$); (from [38]).

•DELPHI 95 [35] (LEP 1)

(See also previous section). The values of E_T of observed jets were larger than 1.5 GeV, while the magnitude of the mass of virtual photon was equal to $< P_1^2 > \approx 0.06 \text{ GeV}^2$, *i.e.* the standard resolved (almost real) photon process, from the point of view of both photons.

Comment: *This is not typical DIS experiment since the photon probe is almost real. The analysis is not typical, either, for the standard large p_T jet study.*

•**DELPHI 97 [39] (LEP 1 and 2)**

Based on the LEP1 and LEP2 data (1995 and 1996 runs), various distributions for the hadronic final state in no-tag events were measured (Fig. 42).




Figure 42: Distributions of: a), d) - invariant mass, b), e) - p_T^{jets} , c) number of reconstructed jets, f) number of events versus pseudorapidity. Solid and dotted lines - VDM + QPM + RPC with GS2 and GRV parton densities, respectively. Dashed line - QCD - RPC contribution for GS2; hatched histogram - VDM part, double hatched - QPM part (from [39]).

Comment: “MC prediction (based on *TWOGAM* generator) gives perfect agreement at CM energy = 91 GeV and slightly exceeds data at 172 GeV”.

•**OPAL 97f [40] (LEP 1.5)**

The inclusive one- and two-jet cross sections have been measured at $\sqrt{s_{ee}} = 130$ and 136 GeV (based on the 1995 run). The anti-tagging condition was applied to one of the initial photons corresponding to maximum photon virtuality $P_{max}^2 \approx 0.8 \text{ GeV}^2$. The cone jet finding algorithm with $R=1$ was used for the first time in photon - photon collisions at LEP.

The transverse energy of jets E_T is taken to be larger than 3 GeV and the pseudorapidity lies within $|\eta| < 1$. Analysis of the transverse energy flow around the jet direction for all jets and in two-jet events was performed.

For the two-photon events, the $x_\gamma^\pm$ distribution and the transverse energy flow around the jet, studied separately for the direct ($x_\gamma > 0.8$) and the double resolved ($x_\gamma < 0.8$) photons contributions, were obtained (see Fig. 43 and Fig. 44, respectively).




Figure 43: The number of two-jet events as a function of x_γ^+ compared to PHOJET (solid line) and PYTHIA (dashed line) simulations. The hatched histogram is the direct contribution to PYTHIA events (from [40]).




Figure 44: The transverse energy flow around the jets in two-jet events. The data are compared with the PHOJET (solid line) and PYTHIA (dashed line) simulations (from [40]).

The one-jet and two-jet cross sections $\frac{d\sigma}{dE_T^{jet}}$ and $\frac{d\sigma}{d\eta^{jet}}$ were measured up to $E_T^{jet} = 16$ GeV, extending the previous measurement at TRISTAN. Results are presented in Fig. 45.




Figure 45: The inclusive one-jet (a) and two-jet (b) cross section as a function of E_T^{jet} for the jets with $|\eta^{jet}| < 1$ compared to the NLO calculations [105]. The solid line is the sum of direct, single - resolved and double - resolved contributions shown separately (from [40]).

The η distributions for one and two jet events were studied as well (see Fig. 46).




Figure 46: The inclusive one-jet (a) and two-jet (b) cross section as a function of E_T^{jet} for the jets with $E_T^{jet} > 3$ GeV compared to the LO QCD calculations of PYTHIA and PHOJET generators (from [40]).

Comment: "The data on η distributions agree well with NLO QCD calculations based on GRV parametrization. The GRV-LO and SaS 1D parametrizations describe the data equally well, the LAC 1, however, gives twice the observed value".

•OPAL 97g [41] (LEP 2)

The dijet production in two photon collisions at e^+e^- energy 161-172 GeV for the average squared virtuality $P^2 = 0.06$ GeV² was measured. The transverse energy of jets is taken to be $E_T > 3$ GeV and the pseudorapidity lies within $|\eta| < 2$. The cone jet finding algorithm with $R=1$ was used.

The direct and resolved photon subprocesses ($x_\gamma^\pm \gtrless 0.8$) were studied. The transverse energy flow around the jet axis is presented in Fig. 47 for the double resolved (a) and direct (b) photon processes. Larger hadronic activity for the resolved photon sample is observed as expected.




Figure 47: The transverse energy flow around the jet axis in two-jet events; a) double-resolved events, (b) direct events (from [41]).

The angular dependence plotted separately for the three basic resolved photon subprocesses and for the direct events is presented in Fig. 48.




Figure 48: The angular distribution of events separated into the direct and the double-resolved contributions, together with the QCD expectations (from [41]).

The cross section for the direct, single- resolved and double-resolved two-jet events versus E_T compared to the NLO calculation with the GRV-HO parton parametrization is shown in Fig. 49.




Figure 49: The inclusive two-jet cross section as a function of E_T^{jet} for jets with $|\eta^{\text{jet}}| < 2$, compared to the NLO calculations [105]. Solid line is the sum of the direct, single-resolved and double-resolved cross sections shown separately (from [41]).

The jet pseudorapidity distribution is presented in Fig. 50 together with the predictions from PYTHIA and PHOJET generators, with various parton parametrizations.




Figure 50: The inclusive two-jet cross sections as a function of $|\eta^{jet}|$ for jets with $E_T^{jet} > 3$ GeV (from [41]).

Comment: "The E_T dependent two-jet cross section is in good agreement with NLO QCD calculation. The GRV-LO and SaS-1D parametrizations describe the inclusive two-jet cross section equally well. The LAC1 parametrization overestimates the inclusive two-jet cross section significantly".

•AMY 92 [42] (TRISTAN)

The measurement of the high p_T hadron production in the quasi-real $\gamma\gamma$ collision was performed (for the energy between 55 and 61.4 GeV). In the observed 3- and 4-jet events there are one or two spectator jets coming from the resolved photon(s). Two parametrizations of parton density in the photon were used: DG and (DO + VDM) (Fig. 51) (The p_T^0 has been introduced to separate the QPM+VDM contributions from the "MJET" one, which corresponds to the RPC contributions).




Figure 51: The experimental p_T^{jet} distribution compared with the predictions of QPM (dotted histogram), QPM + VMD (dashed histogram) and QPM + VMD + MJET (solid histogram) with $p_T^0 = 1.6$ GeV (from [42]).

Comment: *The evidence for hard scattering of hadronic constituents of photons in photon - photon collisions at TRISTAN. "... without the gluonic component it is impossible to reproduce the data".*

•**AMY 1.5 94 [43] (TRISTAN)**

AMY 1.5 (the upgraded AMY detector at the energy 60 GeV) measures the inclusive single- and double-jet cross section $\frac{d\sigma}{dp_T}$ in the range $2.63 \text{ GeV} < p_T < 8.00 \text{ GeV}$ (single jet) and $2.75 \text{ GeV} < p_T < 7.50 \text{ GeV}$ (two jets). The cone algorithm with $R=1$ was used. In Fig. 52 the p_T distributions for one and two jets for $|\eta| < 1.0$, are presented.




Figure 52: The inclusive jet cross section as a function of p_T integrated over $|\eta| < 1.0$ for one-jet (a) and two-jet events (b). The curves represent the sum of QPM (direct) and MJET (resolved) cross sections using the LAC1 (full line), GRV-LO (double dot - dashed), DG (dotted line), LAC3 (dashed line) and LAC1 without the gluon component (dot - dashed line). The short - dashed curve corresponds to the QPM cross section (from [43]).

The η dependence for the jet with p_T larger than 2.5 is presented in Fig. 53.




Figure 53: The inclusive jet cross section as a function of η integrated over $|p_T| \geq 2.5 \text{ GeV}$. The lines convention is the same as in Fig. 52 (from [43]).

Comment: *"The data are in good agreement with LO QCD calculations based on either LAC 1, DG or GRV parametrizations of the parton densities in the photon. The calculation based on LAC 3 disagrees with the data and we do not see the deviation that is observed in the η distribution by the H1 experiment" (here denoted as H1 93).*

•**TOPAZ 93 [44] (TRISTAN)**

The inclusive jet cross section $\frac{d\sigma}{dp_T}$ was measured at the energy of the collision $\sqrt{s_{ee}}=58$ GeV for $< p_T >$ from 2.61 to 7.44 GeV (single jet) and from 2.71 to 7.43 GeV for two-jet events. The cone algorithm with $R = 1$ was used.

The energy flow within the jet as a function of $\delta\eta$ and $\delta\phi$ was also studied and found in agreement with the MC simulation (not shown).

In Figs. 54 and 55 the p_T distribution for the jet and the energy flow per event as a function of the polar angle of the jet are presented, respectively. Fig. 56 shows the inclusive one - jet and two - jet cross sections.




Figure 54: The jet p_T distribution for a no - tag sample. Predictions are shown for the sum of direct + resolved (DG) + VDM processes for different p_T^0 values. Contributions of the direct (dashed line) and VDM (dot - dashed line) processes are shown (from [44]).




Figure 55: The energy flow versus the polar angle θ for the jet sample. The data are compared with the MC predictions for the sum of the direct and the resolved processes (solid histogram) and the direct process only (dashed histogram) (from [44]).




Figure 56: Inclusive a) jet and b) two-jet cross sections as a function of p_T for the pseudorapidity $|\eta_{jet}| \leq 0.7$. TOPAZ Collaboration results from [44].

Comment: The polar angle distribution gives .."direct evidence of the presence of the resolved processes. The data exclude the parametrizations of LAC 3 and DO+ VMD, which predict a very large gluon content even at large x_γ ."

2.3.4 Jet production in the resolved γp scattering

Here we present the data for the jet production in γp scattering taken at HERA collider at $\sqrt{s_{ep}} \sim 300$ GeV¹⁴, where only *one* photon may be resolved¹⁵. The photoproduction events correspond here to the limit of virtuality of the initial photon: $P^2 \sim 4$ GeV² (with median 0.001 GeV²) or (with a special condition) ~ 0.01 -0.02 GeV² (see below). The partonic variables, both related to the initial and final states in the hard partonic subprocesses, are reconstructed from the corresponding quantities for final state hadronic jets. The relation between variables corresponding to these two levels depends on the order of the perturbative QCD calculation, as it was mentioned before.

The parton momentum fraction x_γ^{jet} (called also x_γ^{vis} or x_γ^{obs}) is in the LO case equal to x_γ . In practice it is reconstructed using the two jets with the highest transverse energy E_T^{jet} in the event, using the following relation:

$$x_\gamma^{jet} = \frac{E_T^{jet_1} e^{-\eta^{jet_1}} + E_T^{jet_2} e^{-\eta^{jet_2}}}{2E_\gamma}, \quad (22)$$

where also jet pseudorapidities, $\eta^{jet} = -\ln(\tan(\theta/2))$, and the energy of the photon $E_\gamma (= yE_e)$ enter. (The positive pseudorapidity corresponds to the proton direction.) The scaled energy y of the initial photon (Eq. (4)) is measured from the transverse energy E_T^h and pseudorapidity of hadrons η^h according to the formula

$$y = \frac{1}{2E_e} \sum_h E_T^h e^{-\eta^h}, \quad (23)$$

where the sum is over produced hadrons.

In the following discussion the pseudorapidity of the jet in the laboratory system is denoted by η , whereas η^* means the corresponding variable in the γp CM system. For the HERA collider typically $\eta - \eta^* \sim 2$. The difference of the two jets' transverse energy $E_{T1} - E_{T2}$, ΔE_T , and analogous difference for the pseudorapidities, $\Delta\eta = |\eta_1 - \eta_2|$, are introduced in two-jet events. (For the analysis of the energy of underlying event the variable $\delta\eta = \eta_{cell} - \eta$ is used as in the $\gamma\gamma$ case.) In the analysis of two jet events there appear also the average pseudorapidity $\bar{\eta} = (\eta_1 + \eta_2)/2$ and the average transverse energy of jets, $\bar{E}_T$.

Below, ϕ denotes the azimuthal angle of the particle. In the cone jet algorithm, beside R , also R_{sep} is introduced both in the theoretical calculations and in experimental analysis of the jet production at HERA.

Note also that the multiple interaction included in the experimental analysis is modeled as in the $p\bar{p}$ experiments (see *e.g.* **H1 96a**).

¹⁴ The energy of the electron was at the beginning of running of the HERA collider equal to 26.7 GeV, starting from 1994 it is 27.5 GeV, with the energy of the proton 820 GeV.

¹⁵ The $\gamma\gamma$ events leading to the large p_T jets are rare at HERA.

DATA

•H1 92 [45] (HERA)

The evidence for the hard photoproduction of jet with $E_T > 10$ GeV is reported (the jet cone algorithm with $R < 1$ was used in the analysis).

•H1 93 [46] (HERA)

First measurement of the inclusive jet cross section at the ep collider HERA (based on the 1992 data) is reported. The events with the scaled energy y of the initial photon between 0.25 and 0.7 and the photon virtuality P^2 smaller than 0.01 GeV^2 were collected. The photoproduction of jet was studied for E_T from 7 to 20 GeV and the pseudorapidity interval $-1 < \eta < 1.5$. The jet cone algorithm with $R=1$ (PYTHIA 5.6) was used.

The transverse energy flow around the jet axis was studied and the discrepancy was found in form of too large averaged E_t on the forward side of the jet (see Fig. 57). Among others, multiple parton interactions were mentioned as a possible explanation of this effect.




Figure 57: The transverse energy flow as a function of $\delta\eta$ (integrated over $|\delta\phi| < 1.0$) (a, c) and as a function of $\delta\phi$ (integrated over $|\Delta\eta| < 1.0$) (b, d). Figs. a) and b) correspond to $-1.0 < \eta < 0.5$, b) and c) to $0.5 < \eta < 1.5$ (from [46]).

The inclusive jet cross sections $d\sigma/dE_T$ versus E_T and $d\sigma/d\eta$ as a function of η integrated over the corresponding range of the η and E_T , respectively, were measured and compared with the LO calculation using the following parton parametrizations: for the photon LAC2, LAC3 and GRV-LO, and GRV for the proton (see Fig. 58).




Figure 58: Inclusive jet E_T spectrum integrated over η interval $-1.0 < \eta < 1.5$ (a) and inclusive η spectrum (b) for jets with $E_T > 7$ GeV. LO QCD predictions generated by the PYTHIA generator using parametrizations LAC3 (dashed line), LAC2 (dashed-dotted line), GRV-LO (full line) and GRV-LO (without gluons, dotted line) (from [46]).

The shape of the $d\sigma/dE_T$ is well described in the range of η between -1 and 1.5. It is a problem to describe the η distribution of jets.

Comment: "In the (pseudorapidity) range 0.5 to 1.5 the data show larger average values of the transverse energy flow outside the jet cone on the forward side of the jet than predicted by the MC."

LAC3 gives cross section higher by factor 3 than data for the E_T distribution of the jets. "None of the models describe well the measured η dependence (for jets)."

•H1 95 [47] (HERA)

The photoproduction of 2-jet events (the 1993 data) was studied for the jet E_T range from 7 to 20 GeV and the pseudorapidity interval $0 \leq \eta \leq 2.5$ with $|\Delta\eta| \leq 1.2$ (between the most energetic jets) (see also **H1 96a**). The scaled energy y of the initial photon was between 0.25 and 0.7. Photon virtuality P^2 was smaller than 0.01 GeV^2 . The jet cone finding algorithm with $R=1$ (and 0.7 for cross checks) was used (PYTHIA 5.6 with the GRV-LO for the proton).

For the first time the inclusive (LO) cross sections were derived for the parton level and the gluon density in the photon was measured.

The transverse energy flow around the jet direction per event versus the rapidity distance from the jet direction was studied for $7 \leq E_T \leq 8 \text{ GeV}$ and $0 \leq \eta \leq 1$ and found to be asymmetric and different for the samples with $x_\gamma > 0.4$ and $x_\gamma < 0.4$ (not shown).

The transverse energy flow versus the azimuthal angle around the jet direction and the transverse energy of the underlying events outside the jets (here named E_t^{pedestal} , another name - the underlying event) was studied (see Fig. 59) with the conclusion "the multiple interaction gives an improved description".




Figure 59: (a, c) The transverse energy flow versus the azimuthal angle with respect to the jet direction in two rapidity bins. (b, d) Distributions of transverse energy measured outside of the jets. Histograms show PYTHIA simulations with (full line) and without (dashed line) multiple interactions (from [47]).

To achieve the goal which was here the extraction of the gluon distribution, the single parton cross section $d\sigma/dp_t$ integrated over the parton rapidity range, as well as the single parton $d\sigma/d\eta$ were studied and compared with the LO parametrizations GRV and LAC 1 and LAC 3 (see Fig. 60).




Figure 60: Single parton cross sections: (a) $d\sigma/dp_t$ integrated over the pseudorapidity range $0 < \eta < 2.5$, (b) $d\sigma/d\eta$ for $p_t > 7$ GeV. The solid line - the LO QCD calculation with GRV LO parametrization for partons in the proton and the photon. The dashed (dotted) line - the same for LAC1 (LAC3) parametrizations for the photon (GRV-LO for the proton) (from [47]).

To extract the information on the subprocesses, the full two-jet kinematics was used and the distributions of the $\Delta\eta$, ΔE_T , and of the x_γ and x_p were studied. In Fig. 61 we present the distribution of the x_γ .




Figure 61: The two-jet event distribution of x_γ of the parton from the photon. The solid line - the contribution from the quark resolved photon processes; the dashed line - the direct photon contribution from the PYTHIA MC (from [47]).

The (LO) gluon distribution in the photon was derived, at the average factorization scale $\langle \tilde{Q}^2 \rangle = \langle p_T^2 \rangle = 75 \text{ GeV}^2$ for $0.04 \leq x_\gamma \leq 1$, see the table for numbers:

$\langle \tilde{Q}^2 \rangle$ [GeV ²]	$\langle x_\gamma \rangle$	$x_\gamma g(x_\gamma)/\alpha$ (stat. + syst.)
75	0.059	$1.92 \pm 0.87 \pm 1.68$
	0.14	$1.19 \pm 0.34 \pm 0.59$
	0.33	$0.26 \pm 0.24 \pm 0.33$
	0.59	$-0.12 \pm 0.15 \pm 0.33$
	0.93	$-0.08 \pm 0.61 \pm 0.30$

The results on $x_\gamma g(x_\gamma)/\alpha$ are presented in Fig. 62.




Figure 62: The gluon distribution extracted from the resolved photon processes at $\langle \tilde{Q}^2 \rangle = \langle p_T^2 \rangle = 75 \text{ GeV}^2$. For comparison GRV-LO (full line), LAC1 (dashed) and LAC3 (dotted) gluon parametrizations are shown (from [47]).

Comment: "The multiple interaction option gives an improved description" of the jet profiles and pedestal distributions (still "deviation from data at large jet rapidities $2 < \eta < 2.5$ ").

In the extracting the gluon density q^γ was taken as determined by two-photon experiments at LEP and KEK, in the form given by the GRV-LO parametrization. "A high gluon density at large parton momenta as suggested by the LAC 3 parametrizations is

clearly excluded. The strong rise of the LAC1 parametrization below $x_\gamma \leq 0.08$ is not supported.”

•H1 96a [48] (HERA)

The single jet production with $E_T \geq 7$ GeV (and $-1 < \eta < 2.5$) was measured in ep collisions (data from 199?) with the scaled photon energy $0.25 < y < 0.7$ and P^2 below 0.01 GeV^2 . The cone algorithm with $R=1$ was used.

The properties of the hadronic final state and the distribution of the transverse energy are studied in detail (using PYTHIA, HERWIG and PHOJET generators with GRV-LO parton parametrizations for the proton and the photon). The integration over the γp CM system pseudorapidity $-2.5 < \eta^* < 1$ leads to the total transverse event energy distribution shown in Fig. 63a (here $0.3 < y < 0.7$).

The average transverse energy flow versus η^* for the total E_T range between 25 and 30 GeV was also measured (see Fig. 63b). The shape of both distributions may indicate the need of the multiple interactions.




Figure 63: a) The differential transverse energy cross section integrated over the pseudorapidity ($-2.5 \leq \eta^* \leq 1$). Histograms are the simulations with interactions of the beam remnants (full line - PHOJET, dashed - PYTHIA) and without them (dotted line - PYTHIA). b) The corrected transverse energy flow versus η^* ($\eta^* > 0$ corresponds to the proton direction). The pseudorapidity range and histograms as in a) (from [48]).

To get an insight into the details of the considered events the transverse energy flow outside of the two jets with the highest E_T was studied as a function of x_γ for the $|\eta^*| < 1$ and $\Delta\eta < 1.2$. Results for the transverse energy density can be found in Fig. 64.




Figure 64: The corrected transverse energy density in the region $|\eta^*| < 1$ outside the jets, as a function of x_γ^{jets} . The histograms are as in Fig. 63 (from [48]).

The distribution of the transverse energy around the jet axis was also measured as a function of the $\delta\phi$ (not shown). The jet width obtained in this analysis is similar to the corresponding quantity in the $p\bar{p}$ collision.

Further results obtained for the jet cross section (here $0.25 < y < 0.7$) are presented in Fig. 65a, where $d\sigma/dE_T$ for the jet production in two η regions is shown, and in Fig. 65b, where the distribution $d\sigma/d\eta$ for the events with the transverse jet energy $E_T > 7, 11, 15$ GeV is presented. The comparison with PHOJET and PYTHIA simulations, with and without multiple interactions, was done for both kinds of distributions. Note that the rapidity distribution is more sensitive to the photon structure functions, while the E_T cross section to the matrix elements for the hard processes. Note also that "for E_T bigger than 7 GeV previous measurements (**H1 93**) suffered from a defect and are superseded by this new measurement".

Comment: In addition to the primary hard scattering process, the interaction between the two beam remnants is included in the analysis. It gives "adequate descriptions of data" for the transverse energy versus pseudorapidity and the average energy flow obtained in this analysis. "For the first time the underlying event energy has been measured in jet events using direct and resolved photon probes."

The multiple interaction seems to improve also $d\sigma/dE_T$ and $d\sigma/d\eta$ distributions for jets; within this approach the low E_T and positive η range is still not properly described by the considered two LO parton parametrizations: LAC1 and GRV within PYTHIA program. PHOJET describes these data.




Figure 65: a) Cross section $d\sigma/dE_T$ for the jet production for $E_T^{jet} > 7$ GeV, in two η ranges: $-1 < \eta^{jet} < 2$ and $-1 < \eta^{jet} < 1$. The curves show the MC simulations with interactions of the beam remnants (full line - PHOJET, dashed - PYTHIA) and without them (dotted - PYTHIA). b) Cross section $d\sigma/d\eta^{jet}$ vs η^{jet} for different thresholds in E_T : 7, 11 and 15 GeV. The curves are as in a); additional dash-dotted curve - PYTHIA with LAC1 parametrization (from [48]).

•H1 96b [49] (HERA)

The double differential 2-jet cross section $d\sigma/dx_\gamma^{jets}/d\log(E_T^2/E_0^2)$ was measured as a function of x_γ for different E_T ranges above 8 GeV. Events (the 1994 data) correspond to: P^2 lower than 4 GeV², the scaled energy y between 0.2 and 0.83 and the jet pseudorapidity $-0.5 < \eta < 2.5$, with $\Delta\eta < 1$. The cone algorithm with $R=0.7$ was used. The PYTHIA generator with the multiple interaction using GRV-LO parton parametrizations for the proton and the photon was applied to describe the data.

In Fig. 66 the cross section is shown as a function of the x_γ for the different $\tilde{Q}^2$ ($= E_T^2$) bins (for the E_T^2 distributions - see **H1 97a**).




Figure 66: The double differential two-jet cross section (see the text). The results from the PYTHIA simulation with multiple interactions and GRV-LO parametrizations are shown (from [49]).

For the first time the effective (LL) parton density of the photon

$$\alpha^{-1}x_\gamma(\tilde{q}_\gamma + \frac{9}{4}g_\gamma), \quad (24)$$

with $\tilde{q}_\gamma = \sum(g_\gamma + \bar{q}_\gamma)$, was extracted, for $63 \text{ GeV}^2 < p_T^2 < 1000 \text{ GeV}^2$ and $0.1 < x_\gamma < 0.7$ (see **H1 97a** for the figures).

Comment: "The effective parton distribution grows with the scale p_T^2 , although the increase appears slightly steeper than expected from the GRV-LO parametrization."

•H1 97a [33] (HERA)

A fixed cone algorithm with $R=0.7$ was used to describe jet events with $P^2 < 4 \text{ GeV}^2$ and y between 0.2 and 0.83. The pseudorapidity ranges $0 < \bar{\eta} < 2$ and $\Delta\eta < 1$ as well as $\Delta E_T < 0.5\bar{E}_T$ were selected.

The double differential dijet cross section data as a function of $\bar{E}_T^2$ for few ranges of x_γ are compared with the NLO QCD calculation KK [106] and PYTHIA (GRV) simulation, see Fig. 67.




Figure 67: The double differential cross section as a function of the square of the averaged jet transverse energy and for the different x_γ^{jets} (from [33]).




Figure 68: The effective parton density in the photon (from [33]).

The effective parton distribution in the photon was extracted for $0.2 < x_\gamma < 0.4$ and $0.4 < x_\gamma < 0.7$. Its dependence on the $\tilde{Q}^2$ scale ($= p_T^2$) is shown in Fig. 68.

Comment: "Satisfactory overall description (of the double differential cross section for jet) except for $x_\gamma > 0.6$ ".

•H1 97b [51] (HERA)

The new method of extracting the gluon density in the photon from the charged particles is introduced. Events with $0.3 < y < 0.7$, $P^2 < 0.01 \text{ GeV}^2$ and $|\eta| < 1$ were used. The result on the LO gluon density at $\tilde{Q}^2 = \langle p_T^2 \rangle = 38 \text{ GeV}^2$ is presented in Fig. 69 and compared with the jet data based on the 1993 runs (see also Fig.61).




Figure 69: The gluon density (from [51]).

•**ZEUS 92 [52] (HERA)**

The evidence for the hard scattering in the photoproduction with $E_T > 10$ GeV at HERA is reported (the jet cone algorithm with $R = 1$ was used).

•**ZEUS 94 [53] (HERA)**

The measurement was based on the 1992 data for the single and double jet photoproduction for P^2 below 0.02 GeV^2 for tagged events, otherwise below 4 GeV^2 , and for the y between 0.2 and 0.7. The analysis of the direct and resolved photon processes was made using the HERWIG generator. The jet finding cone algorithm with $R=1$ was used.

The results for the E_T distribution for single jets up to $E_T = 18 \text{ GeV}$, integrated over rapidity η below 1.6, are presented in Fig. 70a. Fig. 70b shows the $d\sigma/d\eta$ data where the disagreement with MC prediction occurs for the positive η .




Figure 70: Inclusive jet distributions for (a) transverse energy of jets, (b) pseudorapidity of jets (from [53]).

Di-jet production has been studied by selecting events with two or more jets with $E_T > 5 \text{ GeV}$, for η smaller than 1.6. (Fig. 71).




Figure 71: Kinematic distributions for events with two or more jets: (a) the jet pair invariant mass, (b) the transverse energy of jets, (c) the pseudorapidity, (d) $\cos \theta^*$ of jet angles in jet-jet CM with respect to the proton momentum for events with $M_{ij} > 16$ GeV. The comparison with MC simulations is shown (from [53]).

The x_γ and x_{proton} distributions were studied as well for events with $|\Delta\eta| < 1.5$, $|\Delta\phi| > 120^\circ$ and the invariant mass of two jets M_{ij} larger than 16 GeV (not shown).

•ZEUS 95a [54] (HERA)

The 1993 data for the production of at least one jet with $E_T > 6$ GeV are presented. Events correspond to P^2 below 4 GeV², y between 0.2 and 0.85 and the jet pseudorapidity between -1 and 2 (-3 and 0 for η^*). PYTHIA 5.6 and HERWIG 5.7 generators (GRV and LAC1 parametrizations for the photon and MRSD₀ for the proton) were used with the cone algorithm for $R=1$. A wider than before range of E_T (up to 41 GeV) and wider η range were considered.

The transverse energy flow around jet axis was studied. Results are presented in Fig. 72, where “there is some discrepancy for the forward-going jets in the $\delta\eta > 1$ ”.




Figure 72: Transverse energy profiles as functions of $\delta\eta$ (top row) and $\delta\phi$ (bottom row) (see the text). Results from the PYTHIA simulation (with both resolved and direct processes) are shown (from [54]).

The E_T distributions integrated over two different pseudorapidity ranges and $d\sigma/d\eta$ distribution integrated above three E_T thresholds: 8, 11 and 17 GeV, are presented in Figs. 73, 74 for different parton parametrizations.

Comment: “In the jet profiles, there is a significant excess of the transverse energy density in the data with respect to the MC expectations for jets in the region $1 < \eta < 2$. This excess is located outside of the jet in the forward direction $\Delta\eta > 1$.”

“Except for the region of very forward, low E_T jets, these measurements are fully consistent with LO QCD predictions in new kinematical regime of the structure of the photon”. The result (for the $d\sigma/d\eta$ for the $E_T > 8$ GeV and the range $-1 < \eta < 1$) does not support the discrepancy of the $d\sigma/d\eta$ with respect to LO QCD calculations observed by the H1 Collaboration (H1 93).”




Figure 73: The E_T distributions for jets (see text). Results of the PYTHIA simulations with LAC1, ACFGP-HO, GS-HO and GRV-HO parametrizations of the parton densities in the photon are shown (from [54]).




Figure 74: The jet pseudorapidity distributions: (a) integrated above three energy thresholds, $E_T > 8, 11$ and 17 GeV (see text); the comparison with PYTHIA simulations using LAC1, GRV-HO parametrizations is shown; (b) only for $E_T > 8$, and compared in addition with ACFGP-HO and GS-HO parametrizations (from [54]).

•ZEUS 95b [55] (HERA)

The photoproduction of dijets, with at least two jets of E_T larger than 6 GeV, is considered in the 1993 data. Events corresponding to the scaled energy y between 0.2 and 0.8 and P^2 lower than 4 GeV² (for $|\Delta\eta| < 0.5$) were grouped in the resolved and direct processes samples. The cone algorithm with $R=1$ was used within the HERWIG

5.7 and PYTHIA 5.6 generators with the GRV-LO parametrization for the photon and the MRSD_L for the proton.

The important x_γ distribution was studied. (The x_γ distribution has also been studied in **ZEUS 96a, 96b, 97b**.) The cut on the x_γ , equal to 0.75, was introduced later to enhance the resolved or the direct photon contributions, and a few distributions were studied separately for these samples.

In Fig. 75 the transverse energy flow around the jet axis versus $\delta\eta$ is shown, for the first time separately for the resolved photon and direct photon contributions (...with the failure to describe low x_γ data).




Figure 75: Transverse energy flow around the jet axis versus $\delta\eta$: (a) $x_\gamma > 0.75$ (direct events); (b) $x_\gamma < 0.75$ (resolved events). Solid (dashed) line - PYTHIA (HERWIG) simulation (from [55]).

The $d\sigma/d\bar{\eta}$ was also measured for the direct and resolved photon events (not shown, see below for comments and new data in **ZEUS 96a**).

Comment: *"Both simulations fail to describe the transverse energy flow in the forward region (see also **ZEUS 95a** and **H1 93**)."*

"The LO QCD predictions (with DG, GRV and GS2 parton parametrizations) lie below the dijet cross section $d\sigma/d\bar{\eta}$ data by factor 1.5-2". The importance of the NLO calculation is stressed.

•ZEUS 96a [56] (HERA)

Analysis of the 1994 data for dijets (2 jets or more) for E_T above 6 GeV and with the jet pair invariant mass above 23 GeV was performed. Events correspond to the range of y between 0.25 and 0.8 and P^2 below 4 GeV². The cone algorithm with $R=1$ was used within PYTHIA 5.7 and HERWIG 5.8 generators (with the MRSA parton parametrization for the proton and the GRV-LO for the photon).

To obtain the scattering angle $\cos\theta^*$ distribution, sensitive to the parton dynamics and not parton densities as in analysis above, the cut not on $\Delta\eta$ (as in previous analysis) but on $\bar{\eta}$ was introduced.

The results for the x_γ , x_p and $\delta\eta$ distributions are shown in Fig. 76. Due to the cut on $\bar{\eta}$ the absolute value of η is restricted to be below 1.8. Note that the applied

cut on the invariant mass suppresses events with low x_γ .

ZEUS 1994




Figure 76: Distributions in x_p (a), x_γ (b) of the transverse energy flow. The resolved ($x_\gamma < 0.75$) and the direct ($x_\gamma > 0.75$) events as a function of $\delta\eta$ are presented separately (c and d, respectively). The MC results are also shown (from [56]).

Very important results concerning the angular distribution due to various partonic subprocesses were obtained for the first time in the large p_T resolved photon processes. The angular distributions $d\sigma/d\cos\theta^*$ for the resolved and direct processes are presented together with the LO and NLO calculation based on the CTEQ3M parametrization for the proton and the GRV for the photon in Fig. 77. The comparison was also made with the HERWIG and PYTHIA simulations (not shown).

Comment: The transverse energy flow is described properly, the " requirements of high mass and small boost remove the disagreement in the forward flow between data and the simulations which has been reported elsewhere in hard photoproduction at HERA."

The dijet angular dependence is well described by the LO and NLO QCD calculations, and also by HERWIG and PYTHIA models.




Figure 77: The dijet angular distributions for the resolved and the direct contributions (see text) (from [56]).

•ZEUS 96b [57] (HERA)

This is the extension of previous analyses (**ZEUS 95, 96a**) based also on the 1994 data on the dijet production with P^2 lower than 4 GeV and y between 0.2 and 0.8. Results for the production of at least two jets with the pseudorapidity between -1.375 and 1.875 and for $E_T^{min} = 6, 8, 11$ and 15 GeV are presented, assuming $|\Delta\eta| < 0.5$. In the data analysis different jet finding algorithms were applied: the cone algorithms EUCELL and PUCELL (both with $R=1$) and the k_T - cluster algorithm KTCLUS.

The resolved cross section was measured in the range $0.3 < x_\gamma < 0.75$ and the direct one - for $x_\gamma > 0.75$. Analysis of the event distribution versus x_γ (see **ZEUS 97b** for the figures), jet profiles in form of the transverse energy flow around the jet axis (presented in Fig. 78) and the $d\sigma/d\bar{\eta}$ for the various jet definition and transverse energy thresholds (see **ZEUS 97a,b** for the new results) were performed.




Figure 78: Jet profiles $\langle dE_T/d\delta\eta \rangle$ in different x_γ bins. Histograms based on the HERWIG generator with (solid line) and without (dashed line) multiparton interactions (from [57]).

Comment: “inclusion of multiparton interactions improves the description (of the jet profiles) significantly.”

•ZEUS 96c [58] (HERA)

The inclusive single jet cross section for the P^2 below 4 GeV^2 with y between 0.2 and 0.8 and for three energy regions for W between 134 and 277 GeV was studied. The measurement of the transverse energy of the jets bigger than 14 GeV and the pseudorapidity range from -1 to 2 was performed. The HERWIG generator (using the jet cone algorithm with $R=1$) including the multiparton interaction was introduced in the analysis. The MRSA parton parametrization was used to describe the proton structure.

Fig. 79 shows the η distributions with the comparison to the LO (PYTHIA) and the NLO QCD predictions (KKS [103]) (Fig. 79 a and b, respectively).




Figure 79: a) The differential cross section $d\sigma/d\eta^{jet}$ integrated over the E_T^{jet} from four thresholds: $E_T^{jet} > 14, 17, 21$ and 25 GeV ; PYTHIA results with the LO parametrizations MRSA for the proton and the GRV-HO and LAC1 for the photon are shown. (b) The same for $E_T^{jet} > 17 \text{ GeV}$ only; curves based on the NLO calculations (KKS [103], using the GRV-HO and the GS-HO parton parametrizations) are displayed (from [58]).

The same is shown (for the NLO approach) in the form of the (data-theory)/theory plot for $E_T > 17 \text{ GeV}$ in Fig. 80.

For events with $E_T > 14 \text{ GeV}$ the η distribution for different $\gamma - p$ CM energy ranges is plotted in Fig. 81.

Comment: “In the region $\eta > 1.5$ the data show a flattening which is not described by the calculations.”




Figure 80: The data as in Fig. 79b with the prediction of the PYTHIA simulation (GRV-HO) and the KKS[103] calculation (GRV-HO and GS-HO parton parametrizations) (from [58]).




Figure 81: The differential cross section $d\sigma/d\eta^{\text{jet}}$ for three regions of the energy W . The PYTHIA results with the MRSA for the proton and with parton distributions in the photon: GRV-HO and LAC1 are shown (from [58]).

•ZEUS 97a [59] (HERA)

The inclusive jet production $d\sigma/d\eta$ was studied with the iterative cone algorithm for events with $P^2 < 4 \text{ GeV}^2$ and $0.2 < y < 0.85$. In the analysis $R_{sep} = R$ or $2R$ was




Figure 82: The differential cross section $d\sigma/d\eta^{jet}$ integrated over E_T^{jet} from four thresholds: $E_T^{jet} > 14, 17, 21$ and 25 GeV ; curves based on the NLO calculations KK[106] using GRV-HO and GS-HO parton parametrizations for the photon and the CTEQ4M for the proton with a) $R=1$, b) $R=0.7$ are shown (from [59]b).

applied. The distributions of jets with $E_T^{min}=14,17,21$ and 25 GeV as a function of η ($-1 < \eta < 2$) were measured. They are not properly described in the forward low E_T region by the NLO calculation KK [106] if $R=1$ is used, for $R=0.7$ the agreement is obtained (Figs. 82 a,b).

The η distributions in three regions of energy W are also in agreement with the NLO calculation for the $R=0.7$, what can be seen in Fig. 83. The data for $R = 1$ (not shown) are not in agreement with a QCD calculation, as in Fig. 81.




Figure 83: The differential cross section $d\sigma/d\eta^{jet}$ for three regions of the energy W . The PYTHIA results with the MRSA for the proton and with the parton distributions in the photon: GRV-HO and LAC1 are shown (from [59]b).

Comment: "The measured cross sections for jets with $R=0.7$ are well described by the (NLL QCD) calculations in the entire range of η ."

•ZEUS 97b [60] (HERA)

The dijet cross section based on the 1994 data for $d\sigma/d\eta$ for jets with $E_T > 6$ GeV, $1.375 < \eta < 1.875$, and with the $|\Delta\eta| < 0.5$ was measured. The different E_T thresholds and the jet definitions (as in **ZEUS 96b**) were implemented. The resolved and the direct photon events ($0.3 < x_\gamma < 0.75$ and $x_\gamma > 0.75$, respectively) were separated in the analysis. The results together with the predictions from a NLO QCD calculation with the additional parameter describing the separation of jets $R_{sep} = R$ or $2R$, as well as corresponding MC simulations (HERWIG 5.8 and PYTHIA 5.7 with or without the multiple interaction) are presented.

The x_γ distribution obtained from the two-jet events indicates a need for the resolved photon contribution (for the first time this kind of measurement was performed

in 1995, see **ZEUS 95b** and **96a,b**). This analysis was based on the KTCLUS algorithm, for the result see Fig. 84. The small x_γ region is not properly described by the MC simulations both with and without the multiple interaction.




Figure 84: The corrected x_γ^{OBS} distribution, solid line - HERWIG with the multiple interaction, dashed line - PYTHIA with the multiple interaction, dotted line - HERWIG without the multiple interaction (from [60]).




Figure 85: The uncorrected transverse energy flow around the jet: solid line - HERWIG with the multiple interaction, dashed line - HERWIG without the multiple interaction (from [60]).

The transverse energy flow obtained using the KTCLUS algorithm is presented for different E_T in Fig. 85, with a similar discrepancy seen in the forward direction, as in previous measurements of the same quantity.

The dijet cross section $d\sigma/d\eta$ obtained under condition $|\Delta\eta| < 0.5$, for different E_T thresholds and with different jet definitions was studied and the results are plotted in Fig. 86a. The predictions of the KK NLO QCD approach [106] are also presented.

The same cross section, now with the KTCLUS jet definition, is plotted in Fig. 86b. The predictions of the KK NLO QCD approach [106] with the different parton parametrizations (GS and GRV) are compared with the data.




Figure 86: a) Dijet cross sections obtained using the different jet algorithms. The curves show the results of the NLO calculation [106] with $R_{sep} = 1$ (solid line) and $R_{sep} = 2$ (dashed) (GS?). b) Dijet cross sections obtained using the definite (KTCLUS) jet algorithm. The curves show the results of the NLO calculation [106] with $R_{sep} = 1$ (solid line) and $R_{sep} = 2$ (dashed) for different parton parametrizations (from [60]).

Comment: The discrepancy in the η distributions for resolved photon contributions was found for events with $E_T^{min} > 6$ GeV, as discussed in ZEUS 96b.

•ZEUS 97c [61] (HERA)

The new 1995 data for the photoproduction of dijets ($P^2 < 4 \text{ GeV}^2$) and y between 0.2 and 0.85 are reported for the first time. The cone algorithm was used in the PYTHIA simulation; both jets have $E_T > 11 \text{ GeV}$ and are in the rapidity range $-1 < \eta < 2$. The preliminary data for $d\sigma/dE_T$ (symmetrized in η ; E_T refers to the highest E_T^{jet}) are shown in Fig. 87.




Figure 87: The dijet cross section $d\sigma/dE_T$ with one jet within the rapidity range $1 < \eta_1 < 2$ and the other jet in three rapidity ranges, described in the figure (from [61]).

The preliminary results for the pseudorapidity distributions are presented in Fig. 88 for the direct and resolved photon samples. Below HERWIG and PYTHIA generators were used with the LL parton parametrizations for the photon. Similar analysis with the comparison to the KK [106] prediction is discussed in **ZEUS 97d**.




Figure 88: The rapidity distribution $d\sigma/d\eta_2$. The extra cut for the leading jet, $E_T > 14 \text{ GeV}$. Three η_1 jet rapidity ranges were considered (as in Fig. 87). The comparison: PYTHIA and HERWIG predictions with the GRV-LO and the GS-LO parametrizations (from [61]).

•ZEUS 97d [62] (HERA)

The k_T - cluster algorithm was used to study the dijet cross sections, $d^3\sigma/dE_T d\eta_1 d\eta_2$, for $E_{T,1} > 14$ GeV and $E_{T,2} > 11$ GeV and the pseudorapidity range $-1 < \eta_1, \eta_2 < 2$ (see also **ZEUS 97c**). Note that y is between 0.2 and 0.85 as before, but $P^2 < 1$ GeV². The E_T dependence was also studied and compared with the KK [106] predictions (with the CTEQ4M and the GS96 parametrization for the proton and the photon, respectively).

Below the results for the E_T distribution in different regions of $\eta_{1,2}$ are presented (Figs. 89 a, b).




Figure 89: a) Dijet cross section $d\sigma/dE_T$ with one jet within the rapidity range $1 < \eta_1 < 2$ and the other jet in three rapidity ranges, described in the figure. b) Dijet cross section $d\sigma/dE_T$ with one jet within rapidity range $0 < \eta_1 < 1$ and the other jet in three rapidity ranges, described in the figure. Comparison with the NLO calculation KK [106] - solid line (dashed line shows the direct contribution) (from [62]).

The rapidity distribution $d\sigma/d\eta_2$ integrated over $E_T > 14$ GeV is presented in Fig. 90. Three η_1 jet rapidity ranges were considered (as in Fig. 88). Here the comparison was made with the NLO calculation by KK [106] with GRV-HO and GS96 parton parametrization for the photon (CTEQ4M parametrization for the proton). The comparison with the HERWIG simulation was also made (not shown, see **ZEUS 97c**).

The high mass ($M_{ij} > 47$ GeV) dijet cross section was also measured. Events with P^2 smaller than 4 GeV² were collected and compared with the same NLO calculation (KK[106]) as above. In this particular analysis the cone algorithm was used with $R=1$. Results are presented in Fig. 91.




Figure 90: The rapidity distribution $d\sigma/d\eta_2$. The extra cut for the leading jet, $E_T > 14$ GeV. Three η_1 jet rapidity ranges were considered (as in Fig. 87). The comparison with the NLO calculation, see text (from [62]).




Figure 91: The dijet distribution $d\sigma/d\cos\theta^*$ for $M_{ij} > 47$ GeV (upper plot) and $d\sigma/dM_{ij}$ for the $|\cos\theta^*| < 0.8$ (lower plot). The comparison with the NLO calculation KK [106] with the CTEQ4M (GS96) parametrization for the proton (photon) is shown (from [62]).

3 Partonic content of the virtual photon

The notion of partonic content of the virtual photon has appeared in high energy interactions soon after the related concept for the real one [1, 13]. Not only the DIS $_{\gamma}$ formalism discussed before can be extended to the scattering on the virtual photon, but also the resolved virtual photon processes can be measured in this case.

Note that for the resolved virtual photon processes the flux (convention dependent) of the virtual photon in the initial electron is introduced in the analysis [63] (see also discussion below). The flux is usually taken (*e.g.* **H1 97**) in a form integrated over the relevant range of y , the scaled energy of the initial photon, and over the squared mass of photon, P^2 :

$$F_{\gamma/e} = \int_{y_{min}}^{y_{max}} dy \int_{P_{min}^2}^{P_{max}^2} dP^2 f_{\gamma/e}(y, P^2) \quad (25)$$

with

$$f_{\gamma/e}(y, P^2) = \frac{\alpha}{2\pi P^2} \left[\frac{1 + (1-y)^2}{y} - \frac{2(1-y)P_{min}^2}{yP^2} \right]. \quad (26)$$

For the photoproduction $P_{min}^2 = m_e^2 y^2 / (1-y)$, where m_e is the electron mass. The assumption of the factorization of the cross section for the ep scattering and for the $\gamma^* p$ process remains valid as long as $p_T^2 \gg P^2$.

3.1 Theoretical framework

The structure function of the virtual photon can be obtained in the Parton Model assuming the production of the $q\bar{q}$ pairs. In the mass parameter range

$$m_q^2 \ll P^2 \ll Q^2 \quad (27)$$

it has a form (to be compared with the Eq. (9)):

$$F_2^{\gamma}(x, Q^2, P^2) = N_c N_f < Q^4 > \frac{\alpha}{\pi} x \{ [x^2 + (1-x)^2] \ln \frac{Q^2}{P^2 x^2} + 6x(1-x) - 2 \},$$

where

$$< Q^4 > = \frac{1}{N_f} \sum_{i=1}^{N_f} Q_i^4. \quad (28)$$

One can see clearly that the scale of the probe has to differ from the P^2 in order to test the structure of the virtual photon (the so called ' P^2 suppression' compared to the real photon case). The QCD evolution equations for the virtual photon are analogous to those for the real photon. Moreover in the case of the virtual photon there is a hope that the initial conditions are not needed, since for $Q^2 \gg P^2 \gg \Lambda_{QCD}^2$ the nonperturbative effects should be absent (see Ref.[13]). The virtual photon may play therefore a unique role in testing the QCD. (The existing parton parametrizations for the virtual photon are shortly discussed in the Appendix.)

The jet production in the processes involving one or two virtual photons can be studied in a similar way as for the real photons. The factorization between the emission of virtual photons and the jet production cross section for $\gamma^*\gamma$, $\gamma^*\gamma^*$ and γ^*p scattering is usually assumed. Note that these cross sections involving virtual initial photons do depend on the convention used for the definition of the flux of virtual photons, since $\sigma^{\gamma^*} \sim 1/\Gamma_{\gamma^*/e}$. At the same time the ee or ep cross sections are free from such ambiguity, as it cancels in the corresponding cross sections according to the (symbolical) relation: $\sigma^e \sim \Gamma_{\gamma^*/e} \sigma^{\gamma^*}$. The notion of the resolved electron may happen to be very useful here (see the next section).

3.2 DIS $_{\gamma^*}$ for virtual photons

Measurements of the virtual photon structure function in the deep inelastic electron scattering are performed using double-tag events. Usually these events were selected in the kinematic region where one of the virtual photons (the probe) has, on the average, a large virtuality Q^2 and the other, the target, a small one, $P^2 \ll Q^2$.

For the DIS $_{\gamma^*}$ measurements, the quantity

$$x_{vis} = Q^2/(Q^2 + P^2 + W_{vis}^2)$$

needs to be converted to the true $x_{Bj} = Q^2/2pq$ variable¹⁶. We start with the old data from PLUTO, the only measurements of “DIS” type for the virtual photon so far.

DATA

•PLUTO 84 [64] (PETRA)

The double - tag events were measured where one of the virtual photons (the probe) had, on the average, virtuality $< Q^2 > = 5 \text{ GeV}^2$ and the other (the target) $< P^2 > = 0.35 \text{ GeV}^2$. (The energy of the beam was here 17.3 GeV.) The experiment was sensitive to the following combination of the virtual photon structure functions:

$$F_{eff} \equiv F_2 + (3/2)F_L.$$

F_{eff} was extracted for x_{Bj} range between 0.05 and 0.6, see Fig. 92. In Fig. 93 the quantity $Q^2\sigma_{\gamma\gamma}/4\pi^2\alpha^2$, averaged over both x_{Bj} and Q^2 , is shown as a function of the measured P^2 (0.2 - 0.8 GeV²).

¹⁶At finite P^2 a modified variable x_{Bj} , which extends over the whole range between 0 and 1 may be introduced. For small ratio $P^2/Q^2 \ll 1$ they coincide.




Figure 92: The PLUTO Collaboration data for the effective structure function for the virtual photon. For the fixed averaged $Q^2 = 5 \text{ GeV}^2$ and $P^2 = 0.35 \text{ GeV}^2$, a dependence on x_{Bj} is shown (from [64]).




Figure 93: The PLUTO Collaboration results for $\frac{Q^2}{4\pi^2\alpha^2} \sigma_{\gamma\gamma}$ for virtual photon as a function of its virtuality P^2 (averaged over x_{Bj} and Q^2 ranges)(from [64]).

3.3 Jet production in resolved virtual photon(s) processes in $\gamma\gamma$ and γp collisions

As for the real photon, the large p_T jets may resolve the virtual photon(s). Provided the corresponding mass relation $\tilde{Q}^2 \sim p_T^2 \gg P_1^2(P_2^2) \gg \Lambda_{QCD}^2$ occurs, one may use the QCD improved parton model as in the case of the real photon (Eqs. (19, 21)).

Recently the LO and the NLO QCD calculations have appeared for the jet production by virtual photon(s) in the considered processes $\gamma\gamma$ and γp .

The transition region between the interaction of an almost real photon and of a virtual photon with the proton is studied by H1 and ZEUS collaborations in the ep collision at HERA. In such analysis the MC generators used to describe the photoproduction and the DIS events as well as the rapidity gap events at HERA are used. As far as the flux of virtual photons is concerned, it is integrated over the corresponding range of the virtuality (see Eqs. (26, 27)).

Some kinematical variables are defined in the γ^*p CM system, and are denoted below by a star, *e.g.* E_T^* . Note that we use the notation P^2 for the squared virtuality of the photon although in the context of the DIS events (on the proton) at HERA it plays the role of the Q^2 .

DATA

•TOPAZ 94 [29] (TRISTAN)

The jet production **TOPAZ 94** data (see discussion in Sec.2.2) are related to the resolved virtual $\gamma(P_1^2)$ +real $\gamma(P_2^2)$ process. The jet production (one and two jets) has been studied with $3.0 \text{ GeV}^2 < P_1^2 < 30 \text{ GeV}^2$, and p_T between 2 and 8 GeV (corresponding to $4 < \tilde{Q}^2 < 64 \text{ GeV}^2$). To what extent one can describe these events using a partonic language for the virtual photon is not clear since in some cases one is probing the region $\tilde{Q}^2/P_1^2$ smaller than 1. (See also Sec.2.2.)

•H1 97 [65] (HERA)

The single jet cross section for the events with $E_T^* > 4\text{-}5 \text{ GeV}$ and with $0.3 < y < 0.8$ is studied in the transition between the photoproduction and the standard DIS_p regime at HERA. The data from the years 1994 and 1995, in three ranges of the squared mass




Figure 94: The inclusive $d\sigma_{ep}/dE_T^*$ jet cross section as a function of the transverse energy E_T^* for various initial photon virtuality P^2 ranges and for the $-2.5 < \eta^* < -0.5$. The HERWIG prediction is denoted by the solid line, the dashed line corresponds to the direct contribution to this model (from [65]).

of the virtual photon $P^2 < 10^{-2} \text{ GeV}^2$, $0.65 < P^2 < 20$ and $9 < P^2 < 49 \text{ GeV}^2$ were collected. The jet pseudorapidity was studied in the range $-2.5 < \eta^* < -0.5$, using the k_T -clustering algorithm and the PHOJET 1.03, LEPTO 6.5 and ARIADNE 4.08, RAPGAP and HERWIG 5.9 generators (with the JETSET used for the hadronization). The GRV 94-HO parton parametrization for the proton and the Drees-Godbole, GRV-HO, SaS-2D parametrizations of the virtual photon were used.

The measured $d\sigma_{ep}/dE_T^*$ as a function of the transverse energy of the jet for various P^2 ranges, integrated over y between 0.3 and 0.6, is presented in Fig. 94. It was found to be in agreement with the HERWIG (DG) model. In Fig. 95 the corresponding data for the rapidity distribution for jets with $E_T^* > 5 \text{ GeV}$ are shown for the various virtuality ranges.




Figure 95: The inclusive $d\sigma_{ep}/d\eta^*$ jet cross section as a function of the rapidity η^* for various initial photon virtuality P^2 ranges for the $E_T^* > 5 \text{ GeV}$. The HERWIG (DG) prediction is denoted by the solid line, the dashed line denotes the direct contribution (from [65]).

The results together with the HERWIG and RAPGAP predictions based on the DG and SaS-2D parton parametrizations in the virtual photon are shown in Fig. 96. To study the dependence of the virtuality of the photon the cross section σ_{γ^*p} is introduced,

$$\sigma_{\gamma^*p \rightarrow \text{jet}+X} = \frac{\sigma_{ep \rightarrow \text{jet}+X}}{F_{\gamma/e}}, \quad (29)$$

although it is not certain that the above factorization really holds for the whole range of kinematical variables.




Figure 96: The inclusive γ^*p jet cross section as a function of $P^2(=Q^2)$ for various ranges of transverse jet energy $E_T^2 = (\vec{Q}^2)$ for the $-2.5 < \eta^* < -0.5$. The HERWIG (DG) prediction is denoted by the dashed line, RAPGAP (DG) - the dotted line, RAPGAP (SaS-2D) - the solid line and the dot-dashed line corresponds to the HERWIG with the GRV-HO parametrization as for the real photon ("no P^2 suppression") (from [65]).

Comment: The HERWIG (DG) model gives "a good description of the data except for jets in the lowest E_t^* range when $9 < Q^2 < 49 \text{ GeV}^2$ ".

•ZEUS 95c [66] (HERA)

The measurement of the direct and resolved photoproduction at HERA with the virtual and the quasi - real photons was performed. In the 1994 run two samples of events were collected: with the photons of virtualities $0.1 \text{ GeV}^2 < P^2 < 0.55 \text{ GeV}^2$ and with the quasi - real photons ($P^2 < 0.02 \text{ GeV}^2$). For each dijet event the fraction of the photon momentum x_γ^{obs} , manifest in the two highest E_T jets ($E_T > 4 \text{ GeV}$), was calculated. The events associated with the direct photon process (high x_γ^{obs}) and with the resolved photon processes (low x_γ^{obs}) were found both in the virtual and quasi - real photon samples. The ratio N_{res}/N_{dir} was calculated as a function of P^2 and it seems to decrease with an increasing photon virtuality, see **ZEUS 97c** for new results.

•ZEUS 97c [67] (HERA)

A study of the ZEUS dijet production data taken during 1995 for the transition region between the photoproduction and the DIS_p is reported. In particular two samples were studied: $0.1 < P^2 < 0.7 \text{ GeV}^2$ and $P^2 < 1.0 \text{ GeV}^2$. The range of the scaled energy was $0.2 < y < 0.55$.




Figure 97: The x_γ distributions for the different squared masses of the initial photon (from [67]).

The jets with $E_T > 6.5$ GeV and $-1.125 < \eta < 1.875$ were used. The HERWIG 5.9 generator with or without the multiple interaction was applied (the MRSA parametrization was used for the proton and the GRV parametrization for the photon). The events corresponding to the direct processes, with $x_\gamma > 0.75$, and to the resolved ones for $x_\gamma < 0.75$, were studied.

The results for the x_γ distribution for the different P^2 ranges are presented in Fig. 97, and the ratio $\sigma_{resolved}/\sigma_{direct}$ versus P^2 is plotted in Fig. 98.




Figure 98: The ZEUS Collaboration results for the ratio of the events associated with the resolved photon to the events due to the direct photon processes as a function of the squared virtuality of the initial photon (from [67]).

4 Related topics

4.1 On leptonic structure functions of the photon

The lepton pair production by two photons in e^+e^- scattering, being one of the basic QED processes,¹⁷ is also a test for experimental methods applied to more complicated two-photon reactions involving hadrons. This applies especially to the process $e^+e^- \rightarrow \text{hadrons}$ where the hadronic structure functions of the photon are measured and the unfolding and tagging methods have to be tested (Secs. 2.1 and 2.2).

The lepton pair production in single tagged events

$$e^+e^- \rightarrow e^+e^-l^+l^- \quad (30)$$

can be described by introducing the QED structure functions for photon: $F_2^{\gamma(QED)}$, $F_1^{\gamma(QED)}$, $F_L^{\gamma(QED)}$, etc (see *e.g.* [68, 69]). Unlike in the hadronic case (*i.e.* with the hadronic final state), these photon structure functions can be reliably calculated in QED. Moreover all the particles in the final state can be directly observed.

Although the final state $e^+e^- \rightarrow e^+e^-e^+e^-$ has been also measured [72], for technical reasons only muonic structure function, denoted below as $F_2^{\gamma(QED)}$, could be extracted (Eq. (30) with $l = \mu$).

$F_L^{\gamma(QED)}$ structure function is much harder to measure, because its contribution is weighted by the small factor y^2 (as for hadronic F_L^γ , see Eq. (3)). However, this longitudinal structure function is not the only structure function that contains additional information. It has been shown that there are azimuthal correlations in the final state particles from two-photon collisions which are sensitive to additional structure functions [70, 71].

If, instead of measuring the cross section $\frac{d\sigma^{e\gamma \rightarrow eX}}{dx_{Bj}dy}$ (Eqs. (2), (3)), one measures in two-body $\gamma^*\gamma$ collision also one final state particle a , additional structure functions F_A^γ and F_B^γ appear [71]:

$$\begin{aligned} \frac{d\sigma(e\gamma \rightarrow eaX)}{dx_{Bj}dyd\Omega_a/4\pi} = & \frac{2\pi\alpha^2}{Q^2} \frac{1 + (1-y)^2}{x_{Bj}y} [(2x_{Bj}\tilde{F}_T^\gamma + \epsilon(y)\tilde{F}_L^\gamma) \\ & - \rho(y)\tilde{F}_A^\gamma \cos\phi_a + \frac{1}{2}\epsilon(y)\tilde{F}_B^\gamma \cos 2\phi_a]. \end{aligned} \quad (31)$$

Here Ω_a describes the direction of particle a in the $\gamma^*\gamma$ rest frame, and ϕ_a is its azimuthal angle around the $\gamma^*\gamma$ axis, relative to the electron (tag) plane. The functions $\epsilon(y)$ and $\rho(y)$ are very close to 1. The standard functions F_T^γ and F_L^γ are obtained from the corresponding $\tilde{F}_i^\gamma$ by integration over the solid angle Ω_a . Note that the formula (31) holds for two leptons or two partons produced in the final state.

The function $F_B^{\gamma(QED)}$ is, in the LL approximation and zero muon mass limit, equal to $F_L^{\gamma(QED)}$, although it involves quite different photon helicity structures. Thus extracting $F_B^{\gamma(QED)}$ can give us indirectly information on $F_L^{\gamma(QED)}$ (in LLA).

¹⁷The earliest experiments on $e^+e^- \rightarrow \text{leptons}$ are summed up in [68, 69].

DATA (Early experiments)

•CELLO 83 [72] (PETRA)

Here both ee and $\mu\mu$ pairs were observed. At $\langle Q^2 \rangle = 9.5 \text{ GeV}^2$ the “muonic” structure function $F_2^{\gamma(QED)}$ was measured¹⁸ and was found to be in good agreement with QED calculations.

•PEP-9/TPC 84 [73] (PEP)

$F_2^{\gamma(QED)}$ was measured at $\langle Q^2 \rangle = 0.3 \text{ GeV}^2$. Results are shown in Fig.105. The quantity $2x_{Bj}F_1^{\gamma(QED)}$ was also extracted in this experiment, from the y region where its contribution was comparable to that of $F_2^{\gamma(QED)}$.

Comment: The importance of $\mu^+\mu^-$ bremsstrahlung background subtraction has been shown in extracting $F_1^{\gamma(QED)}$. Reasonable agreement with QED predictions was obtained, especially for $x_{Bj} \leq 0.25$.

•PLUTO 85 [74] (PETRA)

$F_2^{\gamma(QED)}$ was measured at $\langle Q^2 \rangle = 5.5$ and 40 GeV^2 in the full x_{Bj} range¹⁸. Results are in agreement with QED calculations.

DATA(Recent results)

In the last few years new measurements of the leptonic structure function $F_2^{\gamma(QED)}$ have been performed at LEP by the four collaborations; the additional structure functions $F_{A,B}^{\gamma(QED)}$ have also been measured by ALEPH, L3 and OPAL groups.

Since the invariant mass of the $\mu^+\mu^-$ pair, and hence x_{Bj} , can be determined very accurately, the measurements of $F_2^{\gamma(QED)}$ are only statistically limited. This is in contrast to the hadronic final states, where the extraction of x_{Bj} introduces significant uncertainties. Thus the investigation of QED structure functions is no longer treated only as a test of QED, but rather as a clean experimental procedure meant for testing and refining the experimental procedures to be used in much more complex case of hadronic final states [32].

Because of the precision of LEP data it is also possible to study the effect of the (small) virtuality P^2 of the quasi-real (target) photon.

•ALEPH 97d [75] (LEP 1)

$F_2^{\gamma(QED)}$ has been measured (data from 1994) for $0.6 < Q^2 < 6.3 \text{ GeV}^2$ ($\langle Q^2 \rangle = 2.8 \text{ GeV}^2$) and $3.0 < Q^2 < 60.0 \text{ GeV}^2$ ($\langle Q^2 \rangle = 14.6 \text{ GeV}^2$) (Fig. 99).

¹⁸Results for all $F_2^{\gamma(QED)}$ measurements discussed here will be shown together in Fig. 107.




Figure 99: Structure functions $F_2^{\gamma(QED)}/\alpha$ measured in the ALEPH experiment [75] at $\langle Q^2 \rangle = 2.79 \text{ GeV}^2$, $\langle P^2 \rangle = 0.153 \text{ GeV}^2$ (a) and $\langle Q^2 \rangle = 14.65 \text{ GeV}^2$, $\langle P^2 \rangle = 0.225 \text{ GeV}^2$ (b). (from [75]).

In this experiment also the azimuthal angle distributions have been measured and functions $F_A^{\gamma(QED)}$ and $F_B^{\gamma(QED)}$ extracted (Fig. 100).




Figure 100: The structure functions $F_A^{\gamma(QED)}/F_2^{\gamma(QED)}$ (a) and $\frac{1}{2}F_B^{\gamma(QED)}/F_2^{\gamma(QED)}$ (b) measured in the ALEPH experiment for $\langle Q^2 \rangle = 8.8 \text{ GeV}^2$ [75]; the solid lines are the QED expectations. (from [75]).

Comment: Both $F_2^{\gamma(QED)}$ and $F_A^{\gamma(QED)}$ agree very well with QED predictions. Poorer quality of $F_B^{\gamma(QED)}$ data can be helped with better statistics.

•DELPHI 96a [20] (LEP 1)

$F_2^{\gamma(QED)}$ has been measured at $\langle Q^2 \rangle = 12 \text{ GeV}^2$, as a test for the unfolding and tagging methods in extraction of the hadronic F_2^γ in DELPHI experimental environment (see Sec. 2.1.2). The effect of non - zero target virtuality has been studied (see Fig. 101). A satisfactory fit to the measured $F_2^{\gamma(QED)}$ is obtained for the fixed value of $P^2 = 0.04 \text{ GeV}^2$.




Figure 101: Unfolded $F_2^{\gamma(QED)}/\alpha$ compared with QED predictions obtained for different masses of the target photon: zero mass (upper curve), $\langle P^2 \rangle = 0.04 \text{ GeV}^2$ (middle curve), and average value $\langle P^2 \rangle = 0.13 \text{ GeV}^2$ (lower curve) (from [20]).

•L3 95 [76] (LEP 1)

The data collected by L3 detector in the years 1991-93 were used to extract the $F_2^{\gamma(QED)}$ structure function for $1.4 < Q^2 < 7.6 \text{ GeV}^2$ ($\langle Q^2 \rangle = 1.7 \text{ GeV}^2$) (see Fig. 102). In addition, the angular distribution in the azimuthal angle ϕ_a was measured and some information on previously unmeasured $F_A^{\gamma(QED)}$ and $F_B^{\gamma(QED)}$ was obtained.




Figure 102: $F_2^{\gamma(QED)}/\alpha$ measured at $\langle Q^2 \rangle = 1.7 \text{ GeV}^2$. The effect of nonzero target photon mass is also shown (from [76]).

•L3 97 [77] (LEP 1)

The L3 Collaboration has performed a new measurement of QED photon structure functions: $F_2^{\gamma(QED)}$, $F_A^{\gamma(QED)}$ and $F_B^{\gamma(QED)}$ in the $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$ process [32].

The effect of the photon target virtuality has been studied and is clearly seen in all the three structure functions (see Figs. 103 and 104).




Figure 103: *L3* data on $F_2^{\gamma(QED)}(x, Q^2, P^2)/\alpha$ compared to *QED* predictions at $P^2=0$ (solid line) and $\langle P^2 \rangle_{fit} = 0.033 \text{ GeV}^2$ (dotted line) (from [77]).




Figure 104: *L3* data on structure functions $F_B^{\gamma(QED)}/\alpha$ and $F_A^{\gamma(QED)}/\alpha$ measured at $1.4 < Q^2 < 7.6 \text{ GeV}^2$. The curves are the *QED* predictions for $P^2=0$ (solid) and $\langle P^2 \rangle_{fit} = 0.033 \text{ GeV}^2$ (dashed), respectively (from [77]).

•OPAL 93 [78] (LEP 1)

The *QED* structure function $F_2^{\gamma(QED)}$ for $\langle Q^2 \rangle = 8.0 \text{ GeV}^2$ was extracted from single-tag events at CM energy $\sim M_Z$. In Fig. 105 the data and *QED* expectations are presented together with the earlier measurements at **CELLO 83** [72] and **PEP-9/TPC 84** [73].




Figure 105: $F_2^{\gamma(QED)}/\alpha$ measured at CELLO [72] (open squares), PEP-9 [73] (open circles) and OPAL [78] (full circles). The solid lines are the QED expectations (identical for CELLO and OPAL within the systematic errors) (from [78]).

•OPAL 97h [79] (LEP 1)

Here the extraction of $F_B^{\gamma(QED)}$ was performed for $0.85 < Q^2 < 31 \text{ GeV}^2$ ($\langle Q^2 \rangle = 5.2 \text{ GeV}^2$). $F_A^{\gamma(QED)}$ was not extracted due to the partial integration of the cross section.

The measured value of $\frac{1}{2}\epsilon F_B^{\gamma(QED)}/F_2^{\gamma(QED)}$ is significantly different from zero and its variation with x_{Bj} is consistent with QED (see Fig. 106).




Figure 106: The values of $\frac{1}{2}F_B^{\gamma(QED)}/F_2^{\gamma(QED)}$ obtained from the azimuthal angle distributions (corrected for the effects of the detector). The solid line is the QED prediction for $Q^2=5.2 \text{ GeV}^2$ and $\epsilon=1$ (from [79]).

A compilation of results for the QED structure function $F_2^{\gamma(QED)}$ can be found in Fig. 107 (from [32]). Data from old and new experiments, compared to QED calculations of $F_2^{\gamma(QED)}(x, Q^2, P^2 = 0)$, are presented as a function of x_{Bj} for Q^2 from 0.1 - 40 GeV^2 .




Figure 107: Summary of existing $F_2^{\gamma(QED)}/\alpha$ data for broad Q^2 range shown with QED predictions for $P^2=0$ (from [32]).

4.2 On the structure function of the electron

In the DIS experiments in e^+e^- collisions the inclusive hadron production can be ascribed not to the photon - target but rather to the parent electron or positron - target. This in some cases may be more straightforward than pinning down the structure of the virtual photon, as was mentioned before.

This topic is discussed in *e.g.* Ref. [1]f, [4]. See also the results in [80], where the structure function of the electron (in general - lepton) is related not only to the structure function of photon but also to the electroweak gauge bosons W and Z . The *structure of weak bosons* which appears in this approach was introduced and discussed in Ref. [81].

5 Summary and outlook

The status of recent measurements (~ 1990 and later) of the “structure” of unpolarized real photon in “DIS $_{\gamma}$ ” experiments as well as in large p_T jet production processes involving resolved photon(s) is presented. A qualitative change has appeared in both types of measurements. Final results based on few years’ runs at LEP 1 are being published with the higher statistics and improved unfolding methods. The new data from LEP 1.5 and 2 appeared during last year. On the other hand impressive progress has been obtained in pinning down the individual parton contributions in large p_T resolved photon processes both in e^+e^- and in ep collisions.

The existing discrepancies in describing the final hadronic states in the DIS $_{\gamma}$ experiments as well as in the resolved photon processes in $\gamma\gamma$ and γp collisions seem to have the common origin. The need of additional p_T in the distribution of produced hadrons and jets may suggest an extra interaction involving the constituents of photon(s). One of the possible explanations could be a multiple interaction described in a similar way as introduced in the $p\bar{p}$ processes.

As far as the data related to the “structure” of the virtual photon are concerned, they have just appeared from the hard photon-proton and photon-photon collisions with resolved photon(s). The corresponding new DIS $_{\gamma^*}$ measurement of the structure functions of virtual photon will come probably only with the new generation of accelerators. The interesting extension of the idea to other gauge boson structure functions and the related concept of the structure function of the electron might also be tested there.

The new measurements of the partonic content of the photon are accompanied by the impressive progress made in the NLO QCD calculations for the resolved real and for virtual photon processes.

The future high-energy linear e^+e^- colliders LC as well as related $e\gamma$ and $\gamma\gamma$ colliders, based on the backward Compton scattering on the laser light, will offer a unique opportunity to measure the structure of photon in a new kinematical regime. Moreover, at these colliders measurements of structure functions for photon with a definite polarization should become feasible with a good accuracy (see e.g.[82]).

Acknowledgments.

One of us (MK) would like to thank P. Zerwas, G. Kramer, G. Ingelman, W. Buchmüller for enlightening discussions, and the whole Theory Group for the hospitality during her stay at DESY. She would like to thank Halina Abramowicz and Aharon Levy for figures on F_2^{γ} , important discussions and useful comments. She is also very grateful to Stefan Söldner-Rembold for the compilation of the newest results on F_2^{γ} and to A. Caldwell, M. Erdmann and P.J. Bussey for discussions on HERA results.

We express our special thanks to Aharon Levy for his encouragement, critical reading of the manuscript and very important comments.

We want to thank Zygmunt Ajduk for his help in preparation of this review, we ap-

preciate also fruitful collaboration with Jerzy Rowicki, Jan Żochowski and Paweł Jankowski. We would like to thank Mike Whalley for sending us copies of their "Compilation..." [6].

Supported in part by Polish Committee for Research; grant number 2P03B18209 (1996-1997).

6 Appendix

6.1 Parton parametrizations for the real photon

Duke - Owens (DO) [83]

A leading logarithmic parametrization of the parton distributions in an asymptotic form. Quarks with equal charges have the same distribution functions: $f_u = f_c$, $f_d = f_s$ ($N_f = 4$).

Drees - Grassie (DG) [84]

A parametrization for a full solution of the leading order evolution equations. The input parton distributions with free parameters assumed at $Q_0^2 = 1 \text{ GeV}^2$ and fitted to the only data on F_2^γ existing at that time, at $Q^2 = 5.9 \text{ GeV}^2$, from PLUTO.

Field - Kapusta - Poggioli (FKP) [17]

In this approach F_2^γ is divided into the hadronic part (F_2^{HAD}) and the point-like one (F_2^{PL}). The F_2^{PL} arises from the basic $\gamma^* \rightarrow q\bar{q}$ coupling and higher order QCD corrections, if the final jet p_T is greater than p_T^0 . If p_T is smaller than p_T^0 then the $q\bar{q}$ pair creates a bound state. F_2^{HAD} in this non-perturbative case is taken from the VMD model. The perturbative F_2^{PL} is calculated using the first order splitting functions and the one-loop α_S .

Levy - Abramowicz - Charchuła (LAC) [85]

A parametrization for a full solution of the leading order evolution equation fitted to all available in 1991 measurements of F_2^γ for $Q^2 \geq Q_0^2$. Three sets are provided with the different choices of an input scale Q_0^2 , and the $x \rightarrow 0$ behaviour of a gluon distribution $G(x)$, namely:

- LAC1: $Q_0^2 = 4 \text{ GeV}^2$
- LAC2: $Q_0^2 = 4 \text{ GeV}^2$, $xG(x) \rightarrow \text{const.}$
- LAC3: $Q_0^2 = 1 \text{ GeV}^2$.

Glück - Reya - Vogt (GRV) [86]

The LO and NLO parametrizations of the parton distributions generated dynamically from the valence-like VMD input. The low initial scale $Q_0^2 = 0.3 \text{ GeV}^2$ is universal for the proton, the pion and the photon structure functions. The DIS_γ scheme is introduced to avoid the large- x_{Bj} instability problems. The one free parameter, which is a VMD input normalization, is fixed by the data.

Gordon - Storrow (GS) [87]

The LO and NLO parametrizations. The input structure function at scale $Q_0^2 = 5.3 \text{ GeV}^2$ [87]a and $Q_0^2 = 3 \text{ GeV}^2$ [87]b in the LO analysis is chosen as a sum of a hadronic part from the VMD model and of a point-like part based on the Parton Model. Free parameters (also light quarks masses) are fitted to the data for $Q^2 \geq Q_0^2$. The NLO distributions in the $\overline{\text{MS}}$ are obtained by matching of the F_2 in the LO and the NLO approaches at the Q_0^2 scale.

Aurenche - Chiappetta - Fontannaz - Guillet - Pilon (ACFGP) [88]a

A solution of the NLO evolution equation with the boundary condition taken at $Q_0^2 = 0.25 \text{ GeV}^2$. The input parton distributions was obtained from the VMD model at $Q^2 = 2 \text{ GeV}^2$ and evaluated down to Q_0^2 .

Aurenche - Guillet - Fontannaz [88]b

The NLO parton distributions obtained with the input distributions (shown to be scheme-dependent) at $Q_0^2 = 0.5 \text{ GeV}^2$. The input distributions are based on the VMD model modified to agree with the $\overline{\text{MS}}$ scheme used in this analysis.

Watanabe - Hagiwara - Tanaka - Izubuchi (WHIT) [89]

A set of six the LO parametrizations obtained by fitting the input distributions to all available data for $4 \text{ GeV}^2 \leq Q^2 \leq 100 \text{ GeV}^2$. The parametrizations WHIT1-WHIT6 are based on different input gluon distributions. A massive charm contribution is calculated from the quark parton model and for $Q^2 > 100 \text{ GeV}^2$ from the massive quark evolution equations.

Schuler - Sjöstrand (SaS) [90]

Four sets of the LO parametrizations. The non-perturbative input distributions at $Q_0 = 0.6 \text{ GeV}$ in the SaS1D, SaS1M and $Q_0 = 2 \text{ GeV}$ in the SaS2D, SaS2M sets are based on the VMD model (their normalization is fixed and the x - dependence is obtained from the fits to the data). The fully calculable point-like contribution to the F_2^γ is expressed as an integral of the “state” distributions over the virtuality k^2 of the $\gamma^* \rightarrow q\bar{q}$ state.

The non leading term C_γ is included into the F_2^γ , leading to the $\overline{\text{MS}}$ distributions (SaS1M, SaS2M). $C_\gamma = 0$ gives distributions in the DIS scheme (SaS1D, SaS2D).

6.2 Parton parametrizations for the virtual photon

The Q^2 -dependence of the parton distributions of the virtual photon ($P^2 \neq 0$) for $\Lambda^2 \ll P^2 \ll Q^2$ follows from the corresponding evolution equations [13], as for the real photon ($P^2 = 0$) case. Experimentally important is, however, the low- P^2 region $\Lambda^2 \lesssim P^2$. The parametrizations [1]f, [91, 92], valid for $0 \leq P^2$, are constructed in such a way that they reproduce the distributions of the real photon in the limit $P^2 \rightarrow 0$ and obey the exact Q^2 -evolution equation in the region $\Lambda^2 \ll P^2$.

Schuler - Sjöstrand [92]

An extension of the SaS parton distributions in the real photon to the virtual photon case. In the point-like contribution, the integral over the virtuality of $\gamma^* \rightarrow q\bar{q}$ state, k^2 , is modified by a factor $(\frac{k^2}{k^2+P^2})^2$. In the hadronic contribution a factor $(\frac{m_V^2}{m_V^2+P^2})^2$ is introduced, where m_V is a vector-meson mass.

Drees - Godbole [1]f

Sets of the parton distributions obtained from the corresponding distributions for the real photon (DG) by including multiplicative factors to include properly the $e \rightarrow e\gamma^*$ vertex.

Glück - Reya - Stratmann (GRS) [91]

The LO and NLO distributions obtained by solving the Q^2 -evolution equation with the boundary conditions being a smooth interpolation between the boundary conditions valid at $P^2 = 0$ and for $P^2 \gg \Lambda^2$. The applicability is assumed for the ranges: $P^2 \leq 10 \text{ GeV}^2$, $10^{-4} \leq x \leq 1$, $0.6 \leq Q^2 \leq 5 \cdot 10^4 \text{ GeV}^2$ and $Q^2 \gtrsim 5P^2$.

7 References

- [1] a) V.M. Budnev et al., *Phys. Rep.* C15 (1975) 181
b) H. Kolanoski, *Springer Tracts in Modern Physics* 105 (1984) 187
c) Ch. Berger and W. Wagner, *Phys. Rep.* C146 (1987) 1
d) H. Abramowicz et al., *Int. J. of Modern Physics* A8 (1993) 1005
e) M. Drees and R.M. Godbole, *Pramana - J. of Physics* 41 (1993) 83;
J. of Phys. G: Nucl. and Part. Phys. 21 (1995) 1559
f) M. Drees and R.M. Godbole, *Phys. Rev.* D50 (1994) 3124
g) S. Brodsky and P. Zerwas, *Nucl. Instr. and Methods in Phys. Res.* A355 (1995) 19
h) M. Erdmann, *Springer Tracts in Modern Physics* 138 (1997)
- [2] PLUTO Coll., a) Ch. Berger et al., *Phys. Lett.* B107 (1981) 168; b) Ch. Berger et al., *Phys. Lett.* B142 (1984) 111; c) Ch. Berger et al., *Nucl. Phys.* B281 (1987) 365
CELLO Coll., H.-J. Behrend et al., *Phys. Lett.* B126 (1983) 391
JADE Coll., W. Bartel et al., *Z. Phys.* C24 (1984) 231
TASSO Coll., M. Althoff et al., *Z. Phys.* C31 (1986) 527
TPC/2 γ Coll., H. Aihara et al., *Z. Phys.* C34 (1987) 1; *Phys. Rev. Lett.* 58 (1987) 97
- [3] AMY Coll., T. Sasaki et al., *Phys. Lett.* B252 (1990) 491
- [4] Prakash Mathews and V. Ravindran, *Int. J. of Modern Physics* A11 (1996) 2783
- [5] M. Stratmann and W. Vogelsang, *Z. Phys.* C74 (1997) 641; Proc. "Future physics of HERA" 1995/6, p. 815
- [6] D. Morgan, M.R. Pennington, M.R. Whalley, *J. Phys. G: Nucl. Part. Phys.* 20 (1994) A1
- [7] C.F. von Weizsäcker, *Z. Phys.* 88 (1934) 612; E.J. Williams, *Kgl. Danske Vidensk. Selskab. Mat.-Fiz. Medd.* 13 (1935) N4; E.J. Williams, *Phys. Rev.* 45 (1934) 729
- [8] G. Schuler, Improving the equivalent-photon approximation in electron-positron collisions, CERN-TH/96-297 (hep-ph/9610406)
- [9] V. Telnov, *Nucl. Instr. and Methods in Phys. Res.* A355 (1995) 3
- [10] E. Witten, *Nucl. Phys.* B120 (1977) 189
- [11] R.J. DeWitt et al., *Phys. Rev.* D19 (1979) 2046;
C.H. Llewellyn-Smith, *Phys. Lett.* B79 (1978) 83;
W.R. Frazer, J.F. Gunion, *Phys. Rev.* D20 (1979) 147
- [12] W.A. Bardeen, A.J. Buras, *Phys. Rev.* D20 (1979) 166; *Phys. Rev.* D20 (1980) 2041

- [13] C.T. Hill, G.G. Ross, *Nucl. Phys.* B148 (1979) 373; T. Uematsu, T.F. Walsh *Phys. Lett.* B101 (1981) 263 and *Nucl. Phys.* B199 (1982) 93; G. Rossi, Ph.D.Thesis, UCSD-10p10-227, *Phys. Lett.* B130 (1983) 105
- [14] M. Glück, K. Grassie and E. Reya, *Phys. Rev.* D30, (1984) 1447
- [15] J.J. Sakurai, *Ann. Phys.* 11 (1960) 1; J.J. Sakurai and D. Schildknecht, *Phys. Lett.* B40 (1972) 121
- [16] T.H. Bauer et al., *Rev. of Modern Phys.* 50 (1978) 262 (Err. ibid 51 (1979) 407)
- [17] J.H. Field, F. Kapusta and L. Poggioli, *Phys. Lett.* B181 (1986) 362; J.H. Field, F. Kapusta and L. Poggioli, *Z. Phys.* C36 (1987) 121; F. Kapusta, *Z. Phys.* C42 (1989) 225
- [18] A. Finch (for the ALEPH Coll.), Proc. of the Photon'97 Conference, Egmond aan Zee, The Netherlands, 10-15 May 1997
- [19] ALEPH Coll., submitted to the XVIII Int. Symposium on Lepton-Photon Interactions, Hamburg, July-August 1997 (LP315)
- [20] DELPHI Coll., P. Abreu et al., *Z. Phys.* C69 (1996) 223
- [21] DELPHI Coll., I. Tyapkin, p. 729, Proc. of the 28th Conf. on High Energy Physics ICHEP'96 (Warsaw 1996), eds. Z. Ajduk and A.K. Wróblewski, World Scientific 1997; Kapusta et al., 96-77 Conf 9, Contribution to ICHEP'96
- [22] DELPHI Coll., F. Kapusta et al., submitted to the Int. Europhysics Conference on High Energy Physics, Jerusalem, 1997 (N416, DELPHI 97-122 CONF 104); I. Tyapkin (for the DELPHI Coll.), Proc. of the Photon'97 Conference, Egmond aan Zee, The Netherlands, 10-15 May 1997
- [23] OPAL Coll., R. Akers et al., *Z. Phys.* C61 (1994) 199
- [24] OPAL Coll., K. Ackerstaff et al., *Z. Phys.* C74 (1997) 33
- [25] OPAL Coll., K. Ackerstaff et al., *Phys. Lett.* B412 (1997) 225
- [26] OPAL Coll., K. Ackerstaff et al., *Phys. Lett.* B411 (1997) 387
- [27] AMY Coll., S.K. Sahu et al., *Phys. Lett.* B346 (1995) 208
- [28] AMY Coll., T. Kojima et al., *Phys. Lett.* B400 (1997) 395
- [29] TOPAZ Coll., K. Muramatsu et al., *Phys. Lett.* B322 (1994) 447
- [30] S. Söldner-Rembold, invited talk at the XVIII Int. Symposium on Lepton-Photon Interactions, Hamburg, July-August 1997 (FREIBURG - EHEP - 97 - 19, hep-ex/9711005)

- [31] H. Abramowicz (Plenary talk at ICHEP' 96) p.53, Proc. of the 28th Conf. on High Energy Physics ICHEP'96 (Warsaw 1996), eds. Z. Ajduk and A.K. Wróblewski, World Scientific 1997
- [32] R. Nisius, invited talk at the Int. Europhysics Conference on High Energy Physics, Jerusalem, August 1997 (hep-ex/9712012)
- [33] K. Müller, talk at the Int. Europhysics Conference on High Energy Physics, Jerusalem, August 1997
- [34] ALEPH Coll., submitted to the XVIII Int. Symposium on Lepton-Photon Interactions, Hamburg, July-August 1997 (LP253)
- [35] DELPHI Coll., P. Abreu et al., *Phys. Lett.* B342 (1995) 402
- [36] OPAL Coll., submitted to the XVIII Int. Symposium on Lepton-Photon Interactions, Hamburg, July-August 1997 and to the Int. Europhysics Conference on High Energy Physics, Jerusalem, August 1997 (OPAL Physics Note PN293)
- [37] J.A. Lauber, L. Lönnblad, M.H. Seymour, Proc. of the Photon'97 Conference, Egmond aan Zee, The Netherlands, 10-15 May 1997
- [38] DELPHI Coll., P. Abreu et al., *Z. Phys.* C62 (1994) 357
- [39] N. Zimin (for the DELPHI Coll.), Proc. of the Photon'97 Conference, Egmond aan Zee, The Netherlands, 10-15 May 1997
- [40] OPAL Coll., K. Ackerstaff et al., *Z. Phys.* C73 (1997) 433
- [41] R. Bürgin (for the OPAL Coll.), Proc. of the Photon'97 Conference, Egmond aan Zee, The Netherlands, 10-15 May 1997
- [42] AMY Coll., R. Tanaka et al., *Phys. Lett.* B277 (1992) 215
- [43] AMY Coll., B.J. Kim et al., *Phys. Lett.* B325 (1994) 248
- [44] TOPAZ Coll., H. Hayashii et al., *Phys. Lett.* B314 (1993) 149
- [45] H1 Coll., T. Ahmed et al., *Phys. Lett.* B297 (1992) 205
- [46] H1 Coll., I. Abt et al., *Phys. Lett.* B314 (1993) 436
- [47] H1 Coll., T. Ahmed et al., *Nucl. Phys.* B445 (1995) 195
- [48] H1 Coll., S. Aid et al., *Z. Phys.* C70 (1996) 17
- [49] M. Erdmann for the H1 Coll., contribution to the 28th Conf. on High Energy Physics, Warsaw 1996 (pa02 - 080)
- [50] H1 Coll., C. Adloff et al., *Eur. Phys. J.* C1 (1998) 97

- [51] H1 Coll., submitted to Europhysics Conference on High Energy Physics, Jerusalem, August 1997, N 270
- [52] ZEUS Coll., M. Derrick et al., *Phys. Lett.* B297 (1992) 404
- [53] ZEUS Coll., M. Derrick et al., *Phys. Lett.* B322 (1994) 287
- [54] ZEUS Coll., M. Derrick et al., *Phys. Lett.* B342 (1995) 417
- [55] ZEUS Coll., M. Derrick et al., *Phys. Lett.* B348 (1995) 665
- [56] ZEUS Coll., M. Derrick et al., *Phys. Lett.* B384 (1996) 401
- [57] ZEUS Coll., M. Derrick et al., pa02 - 040, ICHEP' 96, Warsaw 1996
- [58] ZEUS Coll., M. Derrick et al., pa02 - 041, ICHEP' 96, Warsaw 1996
- [59] a. ZEUS Coll., submitted to the Int. Europhysics Conference on High Energy Physics, Jerusalem, August 1997, N650
b. ZEUS Coll., J. Breitweg et al., DESY 98-018
- [60] ZEUS Coll., J. Breitweg, et al., *Eur. Phys. J. C*1 (1998) 109
- [61] M. Hayes (for the ZEUS Coll.), Proc. of the Photon'97 Conference, Egmond aan Zee, The Netherlands, 10-15 May 1997
- [62] ZEUS Coll., submitted to the Int. Europhysics Conference on High Energy Physics, Jerusalem, August 1997, N654
C. Glasman, talk at the Int. Europhysics Conference on High Energy Physics, Jerusalem, August 1997 (hep-ex 9711025)
- [63] F. Halzen, A.D. Martin "Quarks and Leptons: an Introductory Course in Modern Particle Physics", John Wiley & Sons, New York 1984; A. Zembrzuski, M. Krawczyk, "On the Validity of the Equivalent Photon Approximation and the Structure of a Virtual Photon", IFT 18/91, cont. to workshop Physics at HERA 1991, p.617
- [64] PLUTO Coll., Ch. Berger et al., *Phys. Lett.* B142 (1984) 119
- [65] H1 Coll., C. Adloff et al. *Phys. Lett.* B415 (1997) 418;
H. Rick (for the H1 Coll.), Proc. of the Photon'97 Conference, Egmond aan Zee, The Netherlands, 10-15 May 1997
- [66] ZEUS Coll., M. L. Utley, Proc. of the Europhysics Conf. on High Energy Physics, Brussels, 1995 (hep-ex/9508016)
- [67] ZEUS Coll., submitted to the Int. Europhysics Conference on High Energy Physics, Jerusalem, August 1997, N657

- [68] M. Pohl, *Lecture Notes in Physics, Springer Verlag*, 191, p. 234 (1983); W.R. Frazer, A review of two - photon physics, USCD-10P10-231 (Talk presented at the 5th Int. conference on Two-Photon Physics, Aachen, Germany, April, 1983)
- [69] A.C. Caldwell, Talk given at Int. Symposium on Lepton Photon Interactions, Beijing, P.R. China, August, 1995
- [70] N. Arteaga et al., Proc. of the Workshop PHOTON'95, eds. B. Cartwright, D.J. Miller and V.A. Khoze, p.281, World Scientific, Singapore 1995.
- [71] P. Aurenche et al., Physics at LEP2, eds. G. Altarelli, T. Sjöstrand and F. Zwirner, CERN 96-01 (1996) 301
- [72] CELLO Coll., H. - J.Behrend et al., *PL B126* (1983) 384
- [73] PEP-9 Coll, M.P.Cain et al., *Phys. Lett. B147* (1984) 232
- [74] PLUTO COLL. Ch. Berger et al., *Z. Phys. C27* (1985) 249
- [75] C.A. Brew, S. Cartwright, M. Lehto, Proc. of the Photon'97 Conference, Egmond aan Zee, The Netherlands, 10-15 May 1997
- [76] L3 Coll., Contribution to the 1995 Int. Europhysics Conf. on High Energy Physics, Brussels, Belgium, 1995
- [77] L3 Coll., submitted to the Int. Conf. on High Energy Physics, Jerusalem, August 1997
- [78] OPAL Coll., R. Akers et al., *Z. Phys. C60* (1993) 593
- [79] OPAL Coll., K. Ackerstaff et al., *Z. Phys. C74* (1997) 49
- [80] W. Słomiński and J. Szwed, *Acta Physica Polonica* 27 (1996)1887; *Phys. Lett. B387* (1996) 861; hep-ph/9802449
- [81] W. Słomiński and J. Szwed, *Phys. Lett. B323* (1994) 427
- [82] “ e^+e^- collisions at TeV energies: the physics potential”, part D, Proceedings of the Workshop - Annecy, Gran Sasso, Hamburg (1995), DESY 96 - 123D
- [83] D.W. Duke and J.F. Owens, *Phys. Rev. D26* (1982) 1600
- [84] M. Drees and K. Grassie, *Z. Phys. C28* (1985) 451
- [85] H. Abramowicz, K. Charchuła and A. Levy, *Phys. Lett. B269* (1991) 458
- [86] M. Glück, E. Reya and A. Vogt, *Phys. Rev. D46* (1992) 1973
- [87] a. L.E. Gordon and J.K. Storrow, *Z. Phys. C56* (1992) 307
b. L.E. Gordon and J.K. Storrow, *Nucl. Phys. B489* (1997) 405

- [88] a. P. Aurenche, P. Chiapetta, M. Fontannaz, J.P. Guillet, E. Pilon, *Z. Phys.* C56 (1992) 589
b. P. Aurenche, J.P. Guillet and M. Fontannaz, *Z. Phys.* C64 (1994) 621
- [89] K. Hagiwara, M. Tanaka, I. Watanabe and T. Izubuchi, *Phys. Rev.* D51 (1995) 3197
- [90] G.A. Schuler and T. Sjöstrand, *Z. Phys.* C68 (1995) 607
- [91] M. Glück, E. Reya and M. Stratmann *Phys. Rev.* D51 (1995) 3220
- [92] G.A. Schuler, T. Sjöstrand, *Phys. Lett.* B376 (1996) 193
- [93] B.L. Ioffe and A. Oganesian, *Z. Phys.* C69 (1995) 119

Early LO calculations:

- [94] T.-S. Tu, *Nucl. Phys.* B161 (1979) 417; J.F. Owens, *Phys. Rev.* D21 (1980) 54; I. Kang and C.H. Llewellyn Smith, *Nucl. Phys.* B166 (1980) 413; M. Fontannaz et al. *Z. Phys.* C6 (1980) 214
- [95] M. Drees, R. Godbole, *Phys. Rev. Lett.* 61 (1988) 683; *Phys. Rev.* D39 (1989) 169
- [96] H. Baer, J. Ohnemus, J.F. Owens, *Phys. Rev.* D40 (1989) 2844

NLO - one and two jets:

- [97] D. Bödeker, *Phys. Lett.* B292 (1992) 164, *Z. Phys.* C59 (1993) 501
- [98] L.E. Gordon, J.K. Storrow, *Phys. Lett.* B291 (1992) 320
- [99] P. Aurenche, M. Fontannaz, J.P. Guillet, *Phys. Lett.* B338 (1994) 98
- [100] D. Bödeker, G. Kramer, S.G. Salesch, *Z. Phys.* C63 (1994) 471
- [101] M. Greco, A. Vicini, *Nucl. Phys.* B415 (1994) 386
- [102] G. Kramer, S.G. Salesch, *Z. Phys.* C61 (1994) 277
- [103] M. Klasen, G. Kramer, S.G. Salesch, *Z. Phys.* C68 (1995) 113
- [104] P. Aurenche, L. Bourhis, M. Fontannaz, J.P. Guillet, Proc. of "Future Physics at Hera", Hamburg (1996), p.570; ed. G. Ingelman et al.
- [105] T. Kleinwort, G. Kramer, *Nucl. Phys.* B477 (1996) 3; T. Kleinwort, G. Kramer, *Phys. Lett.* B370 (1996) 141
- [106] M. Klasen, G. Kramer, *Phys. Lett.* B366 (1996) 385; *Z. Phys.* C76 (1997) 67; M. Klasen, Ph. Thesis, 1996
- [107] M. Klasen, G. Kramer, *Z. Phys.* C72 (1996) 107

- [108] B.W. Harris, J.F. Owens, presented at the Annual Divisional Meeting of the Division of Particles and Fields of the APS, Minneapolis, USA (1996), hep-ph/9608378
- [109] B.W. Harris, J.F. Owens, *Phys. Rev. D* 56 (1997) 4007; FSU-HEP-971020, hep-ph/9712299
- [110] M. Klasen, T. Kleinwort, G. Kramer, DESY 97-234, ANL-HEP-PK-97-97, hep-ph/9712256

NLO - resolved virtual photon:

- [111] M. Klasen, G. Kramer, B. Pötter, *Eur. Phys. J. C* 1 (1998) 261
- [112] B. Pötter, DESY 97-138, hep-ph/9707319 (1997)